CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 31, 2002

A

Abd-Rabou, S., 1030
Ali, A., 608, 637
Arellano, C., 84
Armstrong, J. W., 523
Auger, P., 648
Aukema, B. H., 1043

B

Backus, E. A., 22, 538
Bancroft, J., 76
Barbercheck, M. E., 84
Barbosa, P., 1097
Barendse, J., 612
Barimo, J. F., 1158
Barrera, J. F., 515
Barret, D., 648
Bates, S. L., 1023
Bazzocchi, G. G., 797
Beerwinkle, K. R., 1
Berberet, R. C., 739
Berger, R., 558
Bernal, J. S., 1219
Bess, E. C., 774
Bextine, B. R., 746
Bisges, A. D., 739
Bitzer, R. J., 449
Blackmer, J. L., 306
Blackwell, S. A., 1168
Blom, P. E., 149, 1110
Boiteau, G., 533
Boivin, G., 1153
Bomgren, S. A., 1231
Booth, S. R., 895
Borchert, D. M., 361
Borden, J. H., 1023
Boulton, T. J., 903
Bounfour, M., 215
Braman, S. K., 564, 1184
Brandenburg, R. L., 476
Brent, C. S., 313
Briano, J. A., 887
Brodeur, J., 177
Brown, J. J., 1037
Brunner, J. F., 995
Bruzzone, O. A., 403
Buckelew, L. D., 449
Buddle, C. M., 438
Bulmer, M. S., 293
Burgio, G., 797
Burnip, G. M., 947
Burns, J. C., 594
Buschman, L. L., 676

C

Calvin, D. D., 864
Cameron, P. J., 65, 367
Cameron, S. J., 215
Cannon, C. A., 977
Cappaert, D. L., 573
Caprio, M. A., 462
Cardona, C., 859
Carlson, J. D., 1062
Carroll, B., 1259
Charlet, L. D., 844
Charlton, R. E., 1142
Chen, C., 215
Chen, F.-Z., 221
Chen, Y., 1037
Chen, Y. H., 626
Chenus, S. A., 482
Chown, S. L., 612
Cladera, J. L., 1009
Clarke, A. R., 941
Cline, A. R., 538
Cloutier, C., 177
Coddington, J. A., 319
Coleman, J. S., 836
Collins, E., 965
Cook, S. P., 668
Cook, T. J., 1091
Corbett, E., 47
Corley, W., 564
Coyle, D. R., 1000
Crist, T. O., 581
Croft, B. A., 648
Cronholm, G. B., 676
Csinos, A. S., 1282
Culbreath, A. K., 1282
Cuperus, G. W., 1062

D

Dawson, T. E., 714
Delamar, Z. D., 1168
Delury, N. C., 947
Devries, T. A., 160
DiFonzo, C. D., 285
Doerr, M. D., 995
Dorn, S., 859
Duan, J. J., 30
Dugravot, S., 550
Dutilleul, P., 54

E

Elliott, N. C., 253
El-Sayed, A. M., 960
Erbilgin, N., 1043
Ergin, E., 168
Erlandson, M., 1135
Eubanks, M. D., 1168, 1175

F

Fei, H., 509
Fell, R. D., 977
Ferreira de Almeida, M. A., 375, 732
Fiacconi, M. R., 797
Fisher, D. S., 594
Fleischer, S. J., 149, 1110
Folgarait, P. J., 403
Fontes, E. M. G., 299
Frank, J. H., 1226
Frouz, J., 608, 637
Funderburk, J., 558

G

Garcia, M. A., 299
Geaghan, J. P., 1
Geden, C. J., 375, 411, 732
Gibb, A. R., 947
Gilbert, L. E., 403
Giles, K. L., 253
Gillespie, D. R., 331
Gillott, C., 1135
Gingras, D., 1153
Ginsberg, H. S., 1191
Gorbet, D., 558
Gould, W. R., 914
Granett, J., 120
Grantham, R. A., 47, 1062
Greenberg, S. M., 142, 397
Griffin, M. L., 107, 112
Guse, C. A., 676
Gutierrez, A. P., 1197

H

Hain, F. P., 668
Hammond, A. M., 1
Han, Q., 462
Han, Y., 602
Hansen, L. D., 1037
Hansen, R. W., 348
Harding, C. L., 1110
Hastings, F. L., 668
Hatle, J. D., 702
Head, G., 30
Heagle, A. S., 594
Heath, J. J., 934
Heimpel, G. E., 1231
Henderson, G., 509
Henderson, I. M., 388
Henneberry, T. J., 306
Herman, R., 753
Herman, R. A., 208

Herms, D. A., 923
Hertl, P. T., 476
Heyer, K., 1191
Higgins, R. A., 676
Hinkle, N. C., 588
Hoffmann, A. A., 965
Hoffmann, M. P., 482
Hofsvang, T., 804
Hogg, D. B., 37
Hollingsworth, R. G., 523
Hollmann, J., 198
Hower, A. A., Jr., 1240
Hoy, C. W., 1251
Hoyt, G. D., 84
Huang, X., 15
Huignard, J., 550
Hull-Sanders, H., 1168
Hummel, R. L., 84
Hung, R. W., 1085
Hunt, J. H., 99

I

Ibaraki, A. I., 198
Isaacs, R., 852

J

Jakobi, V. N., 10
James, S. S., 127
Jarvi, K. J., 160
Jayas, D. S., 816
Jervis, B., 1259
Jian, F., 816
Johnson, D. T., 1259
Johnson, S. J., 1
Jonas, J. L., 1142
Jones, C. G., 836
Jones, R. W., 693
Jones, V. P., 995
Jones, W., 432
Jones, W. A., 397
Jordan, R. A., 1085
Judd, G. J. R., 189
Juliano, S. A., 702

K

Kamijo, K., 1032
Kaplan, I., 1175
Karsai, I., 99
Keena, M. A., 490
Kennedy, A. A., 261
Kennedy, G. G., 84
Kieckhefer, R. W., 253
Kim, D.-S., 686
King, B. H., 505
Klauer, S., 215
Kopper, B. J., 10
Korinus, L. M., 438
Kreiter, S., 648
Krupke, C. H., 189
Kuhar, T. P., 482
Kurota, H., 469

L

Landis, D. A., 285, 880
Lanzoni, A., 797
Lauzière, I., 432
Le Blanc, J.-P. R., 533
Lebrun, R. A., 1191
Lee, J.-H., 686
Lee, L. L., 306
Legaspi, J., 432
Legaspi, J. C., 1219
Legg, D. E., 348
Lewis, B. A., 1259
Lewis, L. C., 1206
Lindgren, B. S., 759, 766
Lindroth, R. L., 10
Linit, M. J., 281
Liu, S.-S., 221
Liu, T.-X., 142, 397
Liu, Y. H., 418
Lloyd, J. E., 348
Lobinske, R. J., 608, 637
Logarzo, G., 914
Lonergan, G., 1268
Lopez, J. D., 1
Loya, L. J., 1240
Luker, L. A., 702
Lundgren, J. G., 1213, 1231
Luo, L., 1
Lykouressis, D. P., 661

M

Mack, T. P., 15
Mallampalli, N., 852
Mandal, B., 1282
Mansfield, S., 723
Manville, J. F., 198
Marshall, D. J., 612
Martin, J. W., 30
Mayer, R. T., 172
McCoy, S., 774
McCraw, B. D., 1259
McCullough, D. G., 261, 584
McCutcheon, G. S., 1030
McGregor, R. R., 331
McLeod, P. J., 1259
McPherson, R. M., 1282
Medina, R. F., 1097
Meng, X.-Z., 602
Mercer, R. D., 612
Michaud, J. P., 827
Michels, G. J., Jr., 253
Miller, D. R., 759, 766
Miller, J. E., 594
Mills, N. J., 723, 1197
Mirkov, T. E., 1219
Molles, M. C., Jr., 774
Momol, T., 558
Monahan, J. F., 668
Morjan, W. E., 1206
Muir, W. E., 816
Mulder, P. G., Jr., 1062, 1259
Mullens, B. A., 588
Munro, V. M. W., 388

N

Nagarkatti, S., 708
Newton, B. L., 92
Nguyen, D., 54
Nguyen, K., 381
Nielsen, K., 1268
Niwa, C. G., 785
Nordheim, E. V., 1043
Nowatzki, T. M., 864

O

Okabayashi, Y., 1052
Omer, A. D., 120
O'Neal, M. E., 285
O'Neil, R. J., 299
Onstad, D. W., 676
Orr, D. B., 361
Osawa, N., 988
Osier, T. L., 10
Otvos, I. S., 903
Overgaard, H. J., 134
Ownley, B. H., 127

P

Parkman, J. P., 1226
Parmenter, R. R., 774
Parrish, C. J., 1168
Peairs, F. B., 676
Peck, R. W., 785
Pedigo, L. P., 449, 1206
Pendley, A. F., 564, 1184
Penny, G. M., 65
Perdikis, D. C., 661
Pereira, R. M., 127
Peruyero, D. B., 693
Pfannenstiel, R. S., 232
Phelan, P. L., 934
Pires, C. S. S., 299
Pires do Prado, A., 375, 732
Pitts, C. W., 273
Polavarapu, S., 1103, 1268
Ponsard, S., 1197
Porter, P., 676
Powell, J. E., 381

R

Raffa, K. F., 1043, 1276
Raghu, S., 941
Ragsdale, D. W., 348
Rau, M. E., 54
Reardon, B. J., 972
Rebek, E. J., 37
Rice, S. A., 47
Richman, D. B., 914
Ring, R. A., 903
Rodriguez, R., 1009
Roitberg, B. D., 189
Roy, M., 177
Rutledge, J., 1276

S
Sahota, T. S., 198
Salvo, A., 874
Sampson, B. J., 339
Sanchez Sanchez, J. A., 331
Sandeson, P. D., 533
Sanon, A., 550
Sappington, T. W., 142, 1128
Sato, H., 1052
Saunders, M. C., 708
Scharff, N., 319
Scherer, P., 753
Scherer, P. N., 208
Schmale, I., 859
Schulze, T. L., 1085
Schuster, D. J., 354
Seal, D. R., 354
Sebolt, D. C., 880
Segura, D. F., 1009
Sétamou, M., 142, 432, 1219
Sewake, K. T., 523
Seymour, R. C., 160
Shanks, C. H., Jr., 895
Sharov, A. A., 1119
Sheng, C.-F., 602
Shimada, M., 469
Shockley, F. W., 22
Simmons, A. M., 1030
Sæthre, M.-G., 804
Skovgård, H., 425
Sloderbeck, P. E., 676
Smilowitz, Z., 149
Smith, H. R., 668
Smith, M. T., 76
Smitley, D. R., 573
Sørensen, L. L., 319
Spence, J. R., 438
Spiers, J. M., 339
Spurgeon, D. W., 754, 972
Stamps, W. T., 281
Stansly, P. A., 354
Stavisky, J., 558
Stephen, F. M., 982
Stewart, C. D., 1184
Stringer, S. J., 339
Strong, W. B., 1023
Suckling, D. M., 947
Sugiura, S., 988
Suh, C. P.-C., 361, 754
Sujii, E. R., 299
Summerville, K. S., 581
Surber, J. B., 30
Sutherst, R. W., 1009
Suwonkerd, W., 134
Szijj, C. E., 588

T
Takagi, M., 134
Tang, Y. Q., 172
Tanigoshi, L. K., 215, 895
Tauber, C. A., 499
Tauber, M. J., 499
Tcheslavskaia, K., 1119
Teixeira, L. A. F., 1103
Thibout, E., 550
Thorpe, K. W., 1119
Thorvilson, H. G., 746
Tipping, C., 558
Tixier, M.-S., 648
Tobin, P. C., 273, 708
Todd, J. W., 1282
Tollefson, J. J., 864
Torres, A. N., 1251
Traniello, J. F. A., 293, 313
Traw, M. B., 714
Trimble, R. M., 960
Tropp, J., 76
Trumble, J. T., 953
Tsai, J. H., 418
Tsuda, Y., 134

U
Uçkan, F., 168
Ulmer, B., 1135

V
Vail, K. M., 127
Valladares, G., 874
Valle, J., 515
Van Duyn, J. W., 361
Vanlaerhoven, S. L., 982
Van Vleet, S. M., 348
Vera, M. T., 1009
Vickerman, D. B., 953
Villa-Castillo, J., 242
Vogel, K., 1268
Vogt, J. T., 47

W
Wäckers, F. L., 859
Wagner, M. R., 242
Wait, D. A., 836
Walgenbach, J. F., 84
Walker, G. P., 65, 367
Walker, M. A., 120
Wallin, K. F., 1276
Wang, C., 381
Watson, J. A., 30
Weathersbee, A. A., III, 172
Wells, M. L., 1282
Welter, S. C., 626
Weseloh, R. M., 1071
Westbrook, J. K., 1, 754
Whiles, M. R., 1142
White, N. D. G., 816
Wicki, M., 1268
Wiedenmann, R. N., 1213
Wigley, P. J., 65
Williams, D. F., 887
Williams, R. N., 934
Williams, T., 515
Wolt, J., 753
Wolt, J. D., 208
Woods, R., 965
Work, T. T., 438, 584
Wright, M. G., 482
Wright, R. E., 47
Wright, R. J., 160

Y
Yeargan, K. V., 92, 107, 112, 232
Young, D. K., 37
Young, D. R., 1158
Young, J. K., 953
Young, L. J., 160
Yuval, B., 941

Z
Zalucki, M. P., 221
Zarrabi, A. A., 739
Zhang, G.-F., 602
Zhioua, E., 1191
Zúñiga, J. A., 515

CUMULATIVE SUBJECT INDEX
Environmental Entomology
Volume 31, 2002

A

abortion,
conelets, 1023
Leptoglossus occidentalis, 1023
Pinus contorta var. *latifolia*, 1023
Pinus monticola, 1023
seed loss, 1023
Abutilon theophrasti, 107, 112
Acalypha ostryaefolia, 107, 112
Coleomegilla maculata, 107, 112
oviposition, 107, 112
predation, 112
whitefly, 107
Zea mays, 107, 112
Acalypha ostryaefolia, 107, 112
Abutilon theophrasti, 107, 112
Coleomegilla maculata, 107, 112
oviposition, 107, 112
predation, 112
whitefly, 107
Zea mays, 107, 112
Acanthoscelides obtectus,
beans, 859
Bruchidae, 859
oviposition behavior, 859
parasitoids, 859
storage pest, 859
Acari,
Azorella selago, 612
community structure, 612
glaciation, 612
habitat specificity, 612
Acheta domesticus,
Hippodamia convergens, 1191
Metarhizium anisopliae, 1191
nontarget effects, 1191
Oncopeltus fasciatus, 1191
pathogenicity, 1191
Acrididae,
barrier island, 1158
coastal vegetation, 1158
grasshoppers, 1158
Orthoptera, 1158
Acrobasis,
blueberry, 852
cherry fruitworm, 852
cranberry fruitworm, 852
Grapholita, 852
Vaccinium, 852
Acrobasis nuxvorella,
models, 1062
pecan, 1062
Pecan nut casebearer, 1062
Aculops lobuliferus,
defoliation, 1000
intensive forestry, 1000
Populus deltoides, 1000
terminal mortality, 1000
adult emergence,
biofix, 864
corn rootworm, 864
Diabrotica, 864
Weibull function, 864
adult longevity,
Cydia pomonella, 804
degree-days, 804
fecundity, 804
lower developmental thresholds, 804
oviposition behavior, 804
adult size,
Apanteles galleriae, 168
fecundity–sex ratio, 168
food quality, 168
immature development, 168
longevity, 168
Aedes aegypti,
development, 54
food availability, 54
mortality, 54
Plagiorchis elegans, 54
time-dependent transition probabilities, 54
age,
flight potential, 1
migration, 1
Pseudaletia unipuncta, 1
sex, 1
tethered-flight, 1
age-specific fecundity,
Anoplophora glabripennis, 76
exotic invasive species, 76
forest pest, 76
intrinsic growth rate, 76
aggregation,
kairomone, 934
Macrodactylus subspinosus, 934
pheromone, 934
plant volatiles, 934
agricultural disturbance,
Collembola, 37
cropping systems, 37
soil ecology, 37
springtails, 37
alfalfa, 92, 253, 953, 1240
aphids, 253
biomagnification, 953
clover root curculio, 1240
Coccinellidae, 253
entomopathogenic nematodes, 1240
Heterorhabditis bacteriophora, 1240
landscape structure, 253
Leiobunum, 92
natural enemies, 253
Phalangium opilio, 92
phenology, 92
phytoextraction, 953
phytoremediation, 953
population, 92
predator-prey interaction, 253
selenium, 953
Sitona hispidulus, 1240
soybean, 92
Spodoptera exigua, 953
Allantonematidae,
biological control, 558
density dependence, 558
Frankliniella fusca, 558
peanut, 558
Thripidae, 558
alley cropping,
Conotrachelus retentus, 281
eastern black walnut, 281
host discrimination, 281
Juglans nigra, 281
resource partitioning, 281
allozyme,
gene flow, 462
genetic structure, 462
Helicoverpa zea, 462
population genetics, 462
allylglucosinolate,
Brassicaceae, 714
glucobrassicin, 714
induced defense, 714
specificity, 714
Amblyomma americanum,
flag sampling, 1085
Ixodes scapularis, 1085
Lyme disease, 1085
Anopheles,
breeding, 134
mosquito larvae, 134
vegetation, 134
water velocity, 134
Anoplophora glabripennis, 76, 490
age-specific fecundity, 76
exotic invasive species, 76
fecundity, 490
forest pest, 76

host effects, 490
intrinsic growth rate, 76
longevity, 490
ant-aphid mutualism,
Aphis gossypii, 1175
Chrysoperla carnea, 1175
Coccinellidae, 1175
predator interference, 1175
Solenopsis invicta, 1175
Anthonomus grandis, 693, 754, 1128
boll weevil, 754, 1128
dioecy, 693
flight behavior, 1128
Hampea, 693
Mexico, 693
pheromone traps, 1128
reproductive ecology, 693
sampling, 1128
supercooling point, 754
wind, 1128
Anthonomus grandis grandis,
boll weevil, 972
critical weight, 972
natural mortality, 972
Apanteles,
Goniozus, 988
mode of parasitism, 988
temporal density-dependence, 988
tortricid leafrollers, 988
Apanteles galleriae,
adult size, 168
fecundity–sex ratio, 168
food quality, 168
immature development, 168
longevity, 168
aphids,
alfalfa, 253
Coccinellidae, 253
landscape structure, 253
natural enemies, 253
predator-prey interaction, 253
Aphis gossypii,
ant-aphid mutualism, 1175
Chrysoperla carnea, 1175
Coccinellidae, 1175
predator interference, 1175
Solenopsis invicta, 1175
apple, 995, 1259
Conotrachelus, 1259
degree-days, 995
development, 995
Lacanobia subjuncta, 995
peach, 1259
phenology model, 995
plum curculio, 1259
sampling, 1259
trap, 1259
Apthona nigriscutis,
biological control, 348
functional lower developmental thresholds, 348
linear degree-day phenology models, 348
Arachis hypogaea,
attractants, 15
carbon dioxide, 15
Elasmopalpus lignosellus, 15
semiochemicals, 15
Araneae,
biodiversity, 438
Carabidae, 438
pitfall trap size, 438
sampling, 438
Staphylindae, 438
Argentina,
biological control, 403
developmental times, 403
fire ant, 403
parasitoid, 403
temperature effects, 403
arthropod community,
disturbance, 774
forest fire, 774
Rio Grande, 774
riparian zone, 774
artificial diet,
Bemisia tabaci, 306
environmental parameters, 306
insect–plant interaction, 306
oviposition, 306
attractants,
Arachis hypogaea, 15
carbon dioxide, 15
Elasmopalpus lignosellus, 15
semiochemicals, 15
attracticide,
behavior, 189
Cydia pomonella, 189
orchard, 189
pheromone, 189
augmentative biological control,
cotton, 361
microclimate, 361
microhabitat, 361
Trichogramma augmentation, 361
Augmentative release,
Chrysoperla rufilabris, 1184
functional response, 1184
Rhinocapsus vanduzeei, 1184
Stephanitis pyrioides, 1184
urban landscape, 1184
azadirachtin,
brown citrus aphid, 172
Lysiphlebus testaceipes, 172
neem extract, 172
Neemix, 172
Toxoptera citricida, 172
Azorella selago,
Acari, 612
community structure, 612
glaciation, 612
habitat specificity, 612

B

Bacillus thuringiensis, 30, 208, 1197, 1213, 1231
beneficial insects, 1213
biological control, 1197, 1231
cabbage, 1231
corn rootworm, 1213
Cry1Ac protein, 30
Cry3Bb, 1213
degradation, 208
environmental risk, 30
enzyme-linked immunosorbent assays, 30
Geocoris punctipes, 1197
insect bioassay, 30
model, 208
Orius tristicolor, 1197
Pieris rapae, 1231
soil, 208
Spodoptera exigua, 1197
strain PS149B1, 208
sugar sprays, 1231
transgenic, 1213
transgenic cotton, 30
Zelus renardii, 1197
Bacillus thuringiensis subsp. *kurstaki*,
Choristoneura occidentalis, 903
Douglas-fir, 903
nontarget Lepidoptera, 903
western spruce budworm, 903
bark beetle,
biological control, 982
directional selection, 1276
heritability, 1276
host acceptance, 1276
plant–insect interactions, 1276
trap catch, 982
vertical distribution, 982
barrier island,
Acrididae, 1158
coastal vegetation, 1158
grasshoppers, 1158
Orthoptera, 1158
Bathyplectes curculionis,
cold tolerance, 739
parasite acclimation, 739
beans,
Acanthoscelides obtectus, 859
Bruchidae, 859
oviposition behavior, 859
parasitoids, 859
storage pest, 859
Beauveria bassiana,
entomopathogenic fungi, 746
microbial control, 746
red imported fire ant, 746
Solenopsis invicta, 746

beet armyworm,
host plants, 142
oviposition preference, 142
Spodoptera exigua, 142
behavior, 189, 476
attracticide, 189
Cydia pomonella, 189
damage, 476
mole crickets, 476
orchard, 189
pheromone, 189
Scapteriscus, 476
soil moisture, 476
surface activity, 476
behavioral ethogram,
electropenetration graph, 538
herbivory, 538
Lygus hesperus, 538
plant bugs, 538
probing behavior, 538
Bemisia tabaci, 306, 1030
artificial diet, 306
biocontrol, 1030
Encarsia, 1030
environmental parameters, 306
Eretmocerus, 1030
insect–plant interaction, 306
oviposition, 306
vegetable, 1030
whitefly, 1030
beneficial insects,
Bacillus thuringiensis, 1213
corn rootworm, 1213
Cry3Bb, 1213
transgenic, 1213
bertha armyworm,
host plant, 1135
Mamestra configurata, 1135
oviposition, 1135
Betula papyrifera,
condensed tannin, 10
Orgyia leucostigma, 10
paper birch, 10
whitemarked tussock moth, 10
binomial sampling,
Dicyphus hesperus, 331
spatial distribution, 331
biocontrol, 388, 1030
Bemisia tabaci, 1030
Encarsia, 1030
Eretmocerus, 1030
foodweb, 388
host competition, 388
leafrollers, 388
shared parasitism, 388
Trigonospila brevifacies, 388
vegetable, 1030
whitefly, 1030
biodiversity, 438, 827
Araneae, 438
biological control, 827
Carabidae, 438
competition, 827
Harmonia axyridis, 827
intraguild predation, 827
multicolored Asian lady beetle, 827
pitfall trap size, 438
sampling, 438
Staphylindae, 438
biofix,
adult emergence, 864
corn rootworm, 864
Diabrotica, 864
Weibull function, 864
biological assessment,
Coleoptera, 1142
indicator species, 1142
Orthoptera, 1142
biological control, 84, 339, 348, 381, 397, 403, 418, 425, 432, 482, 505, 550, 558, 573, 827, 844, 880, 887, 947, 982, 1091, 1168, 1197, 1206, 1219, 1231
Allantonematidae, 558
Apthona nigriscutis, 348
Argentina, 403
Bacillus thuringiensis, 1197, 1231
Bark beetles, 982
biodiversity, 827
Braconidae, 844
brown citrus aphid, 418
cabbage, 1231
Callosobruchus maculatus, 550
Chrysoperla carnea, 1168
Coccinella septempunctata, 1168
codling moth, 947
Coleomegilla maculata, 880
colony size, 1091
competition, 827
conservation tillage, 84
Coptotermes formosanus, 381
Curculionidae, 844
Cydia pomonella, 947
Dasineura oxycoccana, 339
density dependence, 558
development rate, 432
developmental times, 403
Diatraea saccharalis, 1219
Dinarmus basalis, 550
dispersal, 425
dual infections, 887
entomopathogenic nematodes, 381
entomopathogens, 84, 1206
Eoreuma loftini, 432
Eretmocerus, 397
fire ant, 403
fire ants, 1091
Frankliniella fusca, 558
functional lower developmental thresholds, 348
fungicidal effects, 1206
Galerucella calmariensis, 880
Geocoris punctipes, 1197
GNA, 1219
gregarinidae, 573
Harmonia axyridis, 827
herbicide, 1206
Hippodamia convergens, 1168
host age, 505
host size, 505
insecticides, 339
intracolonial prevalence, 887
intraguild predation, 827, 1168
Istocheta, 573
larval behavior, 880
life tables, 482
linear degree-day phenology models, 348
longevity, 432
Lydella jalisco, 432
Lysiphlebia mirzai, 418
mark recapture, 425
microsporidia, 1091
multicolored Asian lady beetle, 827
Musca domestica, 425
nontarget effects, 1206, 1219
Orius tristicolor, 1197
Ostrinia nubilalis, 482
Ovavesicula, 573
parasitoid, 403, 947
parasitoids, 339, 397
peanut, 558
phenology, 947
pheromone, 947
Pieris rapae, 1231
Popillia japonica, 573
red sunflower seed weevil, 844
Reticulitermes flavipes, 381
roundup, 1206
sex ratio, 505
Smicronyx fulvus, 844
snowdrop lectin, 1219
Solenopsis invicta, 887, 1168
Spalangia cameroni, 425, 505
spatial distribution, 425
Spodoptera exigua, 1197
stalkborer, 432
storage system, 550
sugar sprays, 1231
sugarcane, 1219
sulphur-containing compounds, 550
sustainable agriculture, 84
Syrphidae, 339
temperature effects, 403
temperature-dependent development, 418
Thelohania solenopsae, 887

biology,
demography, 354
host, 354
parasitoid, 354
temperature, 354
biomagnification,
alfalfa, 953
phytoextraction, 953
phytoremediation, 953
selenium, 953
Spodoptera exigua, 953
biomonitoring,
developmental instability, 965
environmental stress, 965
Helicoverpa, 965
trait variability, 965
black imported fire ant,
hybrid fire ant, 127
red imported fire ant, 127
Solenopsis invicta, 127
Solenopsis richteri, 127
Thelohania solenopsae, 127
blueberry,
Acrobasis, 852
cherry fruitworm, 852
cranberry fruitworm, 852
Grapholita, 852
Vaccinium, 852
blueberry maggot,
development rates, 1103
diapause, 1103
postdiapause, 1103
Rhagoletis mendax, 1103
body size,
Chrysocharis flacilla, 874
Halticoptera helioponi, 874
Liriomyza huidobrensis, 874
Phaedrotoma scabriventris, 874
tritrophic interactions, 874
boll weevil, 754, 972, 1128
Anthonomus grandis, 754, 1128
Anthonomus grandis grandis, 972
critical weight, 972
flight behavior, 1128
natural mortality, 972
pheromone traps, 1128
sampling, 1128
supercooling point, 754
wind, 1128
brachypterous,
disease management, 1282
Frankliniella fusca, 1282
population dynamics, 1282
tospovirus, 1282
TSWV, 1282
Braconidae,
biological control, 844
Curculionidae, 844
red sunflower seed weevil, 844
Smicronyx fulvus, 844
Triaspis aequoris, 844
Brassica spp.,
mark-recapture, 65
movement, 65
Phthorimaea operculella, 65
Plutella xylostella, 65
potatoes, 65
Brassicaceae,
allylglucosinolate, 714
glucobrassicin, 714
induced defense, 714
specificity, 714
breeding,
Anopheles, 134
mosquito larvae, 134
vegetation, 134
water velocity, 134
brown citrus aphid, 172, 418
azadirachtin, 172
biological control, 418
Lysiphlebia mirzai, 418
Lysiphlebus testaceipes, 172
neem extract, 172
Neemix, 172
temperature-dependent development, 418
Toxoptera citricida, 172
bruchid,
Gleditsia japonica, 469
herbivorous insect, 469
overwintering stage, 469
seasonal adaptation, 469
voltinism, 469
Bruchidae,
Acanthoscelides obtectus, 859
beans, 859
oviposition behavior, 859
parasitoids, 859
storage pest, 859

C

cabbage,
Bacillus thuringiensis, 1231
biological control, 1231
Pieris rapae, 1231
sugar sprays, 1231
California,
Cydia pomonella, 723
nontarget impacts, 723
parasitoid, 723
walnuts, 723
Callosobruchus maculatus,
biological control, 550
Dinarmus basalis, 550
storage system, 550
sulphur-containing compounds, 550
toxicity, 550
Camponotus pennsylvanicus,
carpenter ant, 977
foraging, 977
nutrients, 977
Camponotus vicinus,
habitat, 1037
moisture, 1037
nest aggregation, 1037
nesting site, 1037
temperature, 1037
Carabidae, 242, 438, 785
Araneae, 438
biodiversity, 438
forest health, 242
forest litter arthropods, 785
forest management, 242
Oregon, 785
Pinus ponderosa, 242
pitfall trap size, 438
pitfall traps, 785
prescribed fire, 242, 785
sampling, 438
spiders, 785
Staphylindae, 438
thinning, 242
carbon dioxide,
Arachis hypogaea, 15
attractants, 15
Elasmopalpus lignosellus, 15
semiochemicals, 15
carbon dioxide enrichment,
Tetranychus urticae, 594
Trifolium repens, 594
twospotted spider mite, 594
white clover, 594
carpenter ant,
Camponotus pennsylvanicus, 977
foraging, 977
nutrients, 977
Carposina sasakii,
population dynamics, 686
survivorship, 686
caste,
nutrition, 99
parental manipulation, 99
phenotypic plasticity, 99
Polistes metricus, 99
social wasp, 99
caterpillar predation,
forest, 1097
invertebrate predators, 1097
larval dispersal, 1097
prey size, 1097
cherry fruitworm,
Acrobasis, 852
blueberry, 852
cranberry fruitworm, 852
Grapholita, 852
Vaccinium, 852
China,
Cydia trasias, 602
mass trapping, 602
sex pheromone, 602
Sophora japonica, 602
tortrix, 602

Chironomidae,
Cladotanytarsus, 637
distribution, 637
Glyptotendipes paripes, 637
nuisance midges, 637
population biology, 637
chlorophyll,
fluorescence, 215
raspberry, 215
Tetranychidae, 215
Tetranychus urticae, 215
Choristoneura occidentalis,
Bacillus thuringiensis subsp. *kurstaki*, 903
Douglas-fir, 903
nontarget Lepidoptera, 903
western spruce budworm, 903
Chrysocharis flacilla,
body size, 874
Halticoptera helioponi, 874
Liriomyza huidobrensis, 874
Phaedrotoma scabricentris, 874
tritrophic interactions, 874
Chrysoperla carnea, 1168, 1175
ant-aphid mutualism, 1175
Aphis gossypii, 1175
biological control, 1168
Coccinella septempunctata, 1168
Coccinellidae, 1175
Hippodamia convergens, 1168
intraguild predation, 1168
predator interference, 1175
Solenopsis invicta, 1168, 1175
Chrysoperla rufilabris,
Augmentative release, 1184
functional response, 1184
Rhinocapsus vanduzeei, 1184
Stephanitis pyrioides, 1184
urban landscape, 1184
Cicadellidae,
desert plants, 914
host availability, 914
leafhoppers, 914
Cladotanytarsus,
Chironomidae, 637
distribution, 637
Glyptotendipes paripes, 637
nuisance midges, 637
population biology, 637
Cleridae,
Enoclerus sphegeus, 766
Kairomone, 766
Scolytidae, 766
Thanasimus undatulus, 766
Trypodendron lineatum, 766
climatic factors,
Deois flavopicta, 299
population dynamics, 299
seasonality, 299
survivorship, 299
CLIMEX,
geographical distribution, 1009
medfly, 1009
clover root curculio,
alfalfa, 1240
entomopathogenic nematodes, 1240
Heterorhabditis bacteriophora, 1240
Sitona hispidulus, 1240
coastal vegetation,
Acrididae, 1158
barrier island, 1158
grasshoppers, 1158
Orthoptera, 1158
Coccinella septempunctata,
biological control, 1168
Chrysoperla carnea, 1168
Hippodamia convergens, 1168
intraguild predation, 1168
Solenopsis invicta, 1168
Coccinellidae, 232, 253, 1175
alfalfa, 253
ant-aphid mutualism, 1175
aphids, 253
Aphis gossypii, 1175
Chrysoperla carnea, 1175
diel activity, 232
diurnal, 232
Helicoverpa zea, 232
Hemiptera, 232
landscape structure, 253
natural enemies, 253
predation, 232
predator interference, 1175
predator-prey interaction, 253
Solenopsis invicta, 1175
codling moth, 947, 960
biological control, 947
Cydia pomonella, 947
obliquebanded leafroller, 960
oriental fruit moth, 960
parasitoid, 947
phenology, 947
pheromone, 947, 960
piezoelectric sprayer, 960
tortricids, 960
coffee pest,
Idiarthron subquadratum, 516
mark-recapture, 516
population estimation, 516
sampling, 516
trapping, 516
cold tolerance,
Bathyplectes curculionis, 739
parasite acclimation, 739
Coleomegilla maculata, 107, 112, 880
Abutilon theophrasti, 107, 112
Acalypha ostryaefolia, 107, 112
biological control, 880
Galerucella calmariensis, 880
larval behavior, 880
oviposition, 107, 112
predation, 112
whitefly, 107
Zea mays, 107, 112
Coleoptera,
biological assessment, 1142
indicator species, 1142
Orthoptera, 1142
Collembola, 37, 449
agricultural disturbance, 37
cropping systems, 37
herbicide, 449
microarthropods, 449
population ecology, 449
soil ecology, 37
soil mesofauna, 449
springtails, 37
transgenic soybeans, 449
colony size,
biological control, 1091
fire ants, 1091
microsporidia, 1091
Thelohania solenopsae, 1091
Colorado potato beetle, 149, 1110
geostatistics, 149
global positioning systems, 149
insect spatial dynamics, 149
life table, 1110
population dynamics, 149, 1110
potatoes, 149, 1110
tomatoes, 1110
community structure, 612, 1052
Acari, 612
Azorella selago, 612
function, 1052
glaciation, 612
habitat specificity, 612
leafminers, 1052
parasitoid assemblages, 1052
Phyllonorycter, 1052
Quercus, 1052
competition,
biodiversity, 827
biological control, 827
Harmonia axyridis, 827
intraguild predation, 827
multicolored Asian lady beetle, 827
computer model,
Lymantria dispar, 1071
pathogen, 1071
resting spores, 1071
condensed tannin,
Betula papyrifera, 10
Orgyia leucostigma, 10
paper birch, 10
whitemarked tussock moth, 10
conelets,
abortion, 1023
Leptoglossus occidentalis, 1023

Pinus contorta var. *latifolia*, 1023
Pinus monticola, 1023
seed loss, 1023
conifer,
Ips, 1043
Platysoma, 1043
population dynamics, 1043
predators, 1043
Thanasimus, 1043
connectivity,
Plant structure, 1153
size, 1153
Trichogramma evanescens, 1153
trophic interactions, 1153
Conotrachelus,
apple, 1259
peach, 1259
plum curculio, 1259
sampling, 1259
trap, 1259
Conotrachelus retentus,
alley cropping, 281
eastern black walnut, 281
host discrimination, 281
Juglans nigra, 281
resource partitioning, 281
conservation biological control,
ornamentals, 564
parasitoids, 564
predators, 564
turfgrass, 564
wildflowers, 564
conservation tillage,
biological control, 84
entomopathogens, 84
sustainable agriculture, 84
Coptotermes formosanus, 381, 509
biological control, 381
entomopathogenic nematodes, 381
Reticulitermes flavipes, 381
soldier proportion, 509
temperature, 509
termite survival, 509
wood consumption, 509
corn rootworm, 864, 1213
adult emergence, 864
Bacillus thuringiensis, 1213
beneficial insects, 1213
biofix, 864
Cry3Bb, 1213
Diabrotica, 864
transgenic, 1213
Weibull function, 864
Cotesia glomerata,
Cotesia rubecula, 367
impact, 367
introduction, 367
Pieris rapae, 367
spread, 367
Cotesia rubecula,
Cotesia glomerata, 367
impact, 367
introduction, 367
Pieris rapae, 367
spread, 367
cotton,
augmentative biological control, 361
microclimate, 361
microhabitat, 361
Trichogramma augmentation, 361
cranberry,
Heterorhabditis bacteriophora, 895
Heterorhabditis marelatus, 895
Otiorhynchus sulcatus, 895
Steinernema carpocapsae, 895
strawberry, 895
cranberry fruitworm,
Acrobasis, 852
blueberry, 852
cherry fruitworm, 852
Grapholita, 852
Vaccinium, 852
critical weight,
Anthonomus grandis grandis, 972
boll weevil, 972
natural mortality, 972
crop domestication,
Homoeosoma electellum, 626
native agricultural habitats, 626
parasitism, 626
sunflowers, 626
tritrophic interactions, 626
crop rotation,
oviposition, 285
resistance, 285
cropping systems,
agricultural disturbance, 37
Collembola, 37
soil ecology, 37
springtails, 37
Cry1Ac protein,
Bacillus thuringiensis, 30
environmental risk, 30
enzyme-linked immunosorbent assays, 30
insect bioassay, 30
transgenic cotton, 30
Cry3Bb,
Bacillus thuringiensis, 1213
beneficial insects, 1213
corn rootworm, 1213
transgenic, 1213
Cryptolestes ferrugineus,
mortality, 816
movement, 816
stored-grain, 816
temperature gradient, 816
Curculionidae,
biological control, 844
Braconidae, 844
red sunflower seed weevil, 844
Smicronyx fulvus, 844
Triaspis aequoris, 844
Cydia pomonella, 189, 723, 804, 947
adult longevity, 804
attracticide, 189
behavior, 189
biological control, 947
California, 723
codling moth, 947
degree-days, 804
fecundity, 804
lower developmental thresholds, 804
nontarget impacts, 723
orchard, 189
oviposition behavior, 804
parasitoid, 723, 947
phenology, 947
pheromone, 189, 947
walnuts, 723
Cydia trasias,
China, 602
mass trapping, 602
sex pheromone, 602
Sophora japonica, 602
tortrix, 602

D

damage,
behavior, 476
mole crickets, 476
Scapteriscus, 476
soil moisture, 476
surface activity, 476
Dasineura oxycoccana,
biological control, 339
insecticides, 339
parasitoids, 339
Syrphidae, 339
Vaccinium, 339
decision rules,
Listronotus oregonensis, 1251
Petroselinum crispum, 1251
probability density functions, 1251
spatial pattern, 1251
transect pattern, 1251
defoliation,
Aculops lobuliferus, 1000
intensive forestry, 1000
Populus deltoides, 1000
terminal mortality, 1000
degradation,
Bacillus thuringiensis, 208
model, 208

soil, 208
strain PS149B1, 208
degree-day models,
diapause, 708
Endopiza viteana, 708
grape berry moth, 708
phenology, 708
degree-days, 804, 995
adult longevity, 804
apple, 995
Cydia pomonella, 804
development, 995
fecundity, 804
Lacanobia subjuncta, 995
lower developmental thresholds, 804
oviposition behavior, 804
phenology model, 995
demography, 120, 354
biology, 354
grape phylloxera, 120
host, 354
parasitoid, 354
plant growth stage, 120
population growth, 120
temperature, 354
dendrobium orchids,
Frankliniella occidentalis, 523
scouting methods, 523
thrips, 523
Dendroctonus ponderosae,
Hylastes longicollis, 759
Hylurgops porosus, 759
Ips latidens, 759
Ips pini, 759
Pinus contorta, 759
density,
dispersal, 533
flight, 533
Leptinotarsa decemlineata, 533
density dependence,
Allantonematidae, 558
biological control, 558
Frankliniella fusca, 558
peanut, 558
Thripidae, 558
Deois flavopicta,
climatic factors, 299
population dynamics, 299
seasonality, 299
survivorship, 299
desert plants,
Cicadellidae, 914
host availability, 914
leafhoppers, 914
development, 54, 588, 608, 797, 995
Aedes aegypti, 54
apple, 995
degree-days, 995
Diptera, 588
Fannia, 588
food availability, 54
Glyptotendipes paripes, 608
Lacanobia subjuncta, 995
leafminers, 797
life stages, 608
Liriomyza huidobrensis, 797
Liriomyza trifolii, 797
manure, 588
moisture, 588
mortality, 54, 797
nuisance midge, 608
phenology model, 995
Plagiorchis elegans, 54
temperature, 608
temperature dependence, 797
time-dependent transition probabilities, 54
development rate, 375, 432
biological control, 432
blueberry maggot, 1103
diapause, 1103
Eoreuma loftini, 432
longevity, 375, 432
Lydella jalisco, 432
muscoid flies, 375
postdiapause, 1103
Rhagoletis mendax, 1103
stalkborer, 432
Tachinaephagus zealandicus, 375
temperature, 375
development time,
Plutella xylostella, 221
rate function, 221
survival, 221
temperature, 221
developmental instability,
biomonitoring, 965
environmental stress, 965
Helicoverpa, 965
trait variability, 965
developmental rate,
nonlinear models, 177
predator-prey synchrony, 177
Stethorus punctillum, 177
temperature, 177
Tetranychus mcdanieli, 177
developmental temperature threshold,
egg development, 661
Macrolophus pygmaeus, 661
Miridae, 661
nymphal development, 661
thermal constant, 661
developmental times,
Argentina, 403
biological control, 403
fire ant, 403
parasitoid, 403
temperature effects, 403
Diabrotica,
adult emergence, 864
biofix, 864
corn rootworm, 864
Weibull function, 864
diapause, 708, 1103
blueberry maggot, 1103
degree-day models, 708
development rates, 1103
Endopiza viteana, 708
grape berry moth, 708
phenology, 708
postdiapause, 1103
Rhagoletis mendax, 1103
Diatraea grandiosella,
European corn borer, 676
Ostrinia nubilalis, 676
resistance management, 676
simulation modeling, 676
southwestern corn borer, 676
Diatraea saccharalis,
biological control, 1219
GNA, 1219
nontarget effects, 1219
snowdrop lectin, 1219
sugarcane, 1219
Dicyphus hesperus,
binomial sampling, 331
spatial distribution, 331
diel activity,
Coccinellidae, 232
diurnal, 232
Helicoverpa zea, 232
Hemiptera, 232
predation, 232
diet,
foraging, 47
liquid food, 47
Solenopsis invicta, 47
solid food, 47
success rates, 47
Dihrinus himalayanus,
Musca domestica, 411
Muscidifurax raptor, 411
Spalangia cameroni, 411
Spalangia endius, 411
Spalangia gemina, 411
Dinarmus basalis,
biological control, 550
Callosobruchus maculatus, 550
storage system, 550
sulphur-containing compounds, 550
toxicity, 550
dioecy,
Anthonomus grandis, 693
Hampea, 693
Mexico, 693
reproductive ecology, 693
Diptera,
development, 588
Fannia, 588

manure, 588
moisture, 588
directional selection,
bark beetle, 1276
heritability, 1276
host acceptance, 1276
plant–insect interactions, 1276
disease management,
brachypterous, 1282
Frankliniella fusca, 1282
population dynamics, 1282
tospovirus, 1282
TSWV, 1282
disparlure,
gypsy moth, 1119
Lymantria dispar, 1119
mating disruption, 1119
pheromone, 1119
release-recapture, 1119
dispersal, 425, 533
biological control, 425
density, 533
flight, 533
Leptinotarsa decemlineata, 533
mark recapture, 425
Musca domestica, 425
Spalangia cameroni, 425
spatial distribution, 425
distribution,
Chironomidae, 637
Cladotanytarsus, 637
Glyptotendipes paripes, 637
nuisance midges, 637
population biology, 637
disturbance,
arthropod community, 774
forest fire, 774
Rio Grande, 774
riparian zone, 774
diurnal,
Coccinellidae, 232
diel activity, 232
Helicoverpa zea, 232
Hemiptera, 232
predation, 232
domatia,
Kampimodromus aberrans, 648
leaf trichomes, 648
pollen, 648
vineyards, 648
Douglas-fir,
Bacillus thuringiensis subsp. *kurstaki*, 903
Choristoneura occidentalis, 903
nontarget Lepidoptera, 903
western spruce budworm, 903
dual infections,
biological control, 887
intracolonial prevalence, 887
Solenopsis invicta, 887
Thelohania solenopsae, 887
Vairimorpha invictae, 887

E

eastern black walnut,
alley cropping, 281
Conotrachelus retentus, 281
host discrimination, 281
Juglans nigra, 281
resource partitioning, 281
egg development,
developmental temperature threshold, 661
Macrolophus pygmaeus, 661
Miridae, 661
nymphal development, 661
thermal constant, 661
Elasmopalpus lignosellus,
Arachis hypogaea, 15
attractants, 15
carbon dioxide, 15
semiochemicals, 15
electropenetration graph,
behavioral ethogram, 538
herbivory, 538
Lygus hesperus, 538
plant bugs, 538
probing behavior, 538
Empoasca fabae,
exudates, 22
host plant resistance, 22
Medicago sativa, 22
plant volatiles, 22
Encarsia,
Bemisia tabaci, 1030
biocontrol, 1030
Eretmocerus, 1030
vegetable, 1030
whitefly, 1030
Endopiza viteana,
degree-day models, 708
diapause, 708
grape berry moth, 708
phenology, 708
energetic reserves,
kairomones, 941
nutrition, 941
pharmacophagy, 941
pheromones, 941
survival, 941
Enoclerus sphegeus,
Cleridae, 766
Kairomone, 766
Scolytidae, 766
Thanasimus undatulus, 766
Trypodendron lineatum, 766
entomopathogenic fungi,
Beauveria bassiana, 746
microbial control, 746
red imported fire ant, 746
Solenopsis invicta, 746
entomopathogenic nematode,
multiparasitism, 1226
Ormia depleta, 1226
Scapteriscus spp., 1226
Steinernema scapterisci, 1226
tachinid, 1226
entomopathogenic nematodes, 381, 1240
alfalfa, 1240
biological control, 381
clover root curculio, 1240
Coptotermes formosanus, 381
Heterorhabditis bacteriophora, 1240
Reticulitermes flavipes, 381
Sitona hispidulus, 1240
entomopathogens, 84, 1206
biological control, 84, 1206
conservation tillage, 84
fungicidal effects, 1206
herbicide, 1206
nontarget effects, 1206
roundup, 1206
sustainable agriculture, 84
environmental parameters,
artificial diet, 306
Bemisia tabaci, 306
insect–plant interaction, 306
oviposition, 306
environmental risk,
Bacillus thuringiensis, 30
Cry1Ac protein, 30
enzyme-linked immunosorbent assays, 30
insect bioassay, 30
transgenic cotton, 30
environmental stress,
biomonitoring, 965
developmental instability, 965
Helicoverpa, 965
trait variability, 965
enzyme-linked immunosorbent assays,
Bacillus thuringiensis, 30
Cry1Ac protein, 30
environmental risk, 30
insect bioassay, 30
transgenic cotton, 30
Eoreuma loftini,
biological control, 432
development rate, 432
longevity, 432
Lydella jalisco, 432
stalkborer, 432
Eretmocerus, 397, 1030
Bemisia tabaci, 1030
biocontrol, 1030
biological control, 397
Encarsia, 1030
parasitoids, 397
tomato, 397
tritrophic, 397
vegetable, 1030
whitefly, 397, 1030

European corn borer, 160, 676
Diatraea grandiosella, 676
geostatistics, 160
maize, 160
Ostrinia nubilalis, 676
resistance management, 676
simulation modeling, 676
southwestern corn borer, 676
Exomala orientalis,
highbush blueberries, 1268
mating disruption, 1268
ornamental nurseries, 1268
sex pheromone, 1268
(Z)-7-tetradecen-2-one, 1268
exotic invasive species,
age-specific fecundity, 76
Anoplophora glabripennis, 76
forest pest, 76
intrinsic growth rate, 76
exotic pest,
Ips pini, 261
nonindigenous insect, 261
pheromone trapping, 261
predators, 261
Thanasimus dubius, 261
exudates,
Empoasca fabae, 22
host plant resistance, 22
Medicago sativa, 22
plant volatiles, 22

F

Fannia,
development, 588
Diptera, 588
manure, 588
moisture, 588
fecundity, 313, 490, 804
adult longevity, 804
Anoplophora glabripennis, 490
Cydia pomonella, 804
degree-days, 804
host effects, 490
longevity, 490
lower developmental thresholds, 804
nitrogen limitation, 313
ovaries, 313
oviposition behavior, 804
reproduction, 313
termites, 313
Zootermopsis angusticollis, 313
fecundity–sex ratio,
adult size, 168
Apanteles galleriae, 168
food quality, 168
immature development, 168
longevity, 168
feeding treatment,
host density, 732
muscoid flies, 732
parasitoid female age, 732
Tachinaephagus zealandicus, 732
temperature, 732
fertilization,
growth/differentiation balance, 923
host plant resistance, 923
secondary metabolism, 923
woody plants, 923
fire ant,
Argentina, 403
biological control, 403, 1091
colony size, 1091
developmental times, 403
microsporidia, 1091
parasitoid, 403
temperature effects, 403
Thelohania solenopsae, 1091
flag sampling,
Amblyomma americanum, 1085
Ixodes scapularis, 1085
Lyme disease, 1085
flight,
density, 533
dispersal, 533
Leptinotarsa decemlineata, 533
flight behavior,
Anthonomus grandis, 1128
boll weevil, 1128
pheromone traps, 1128
sampling, 1128
wind, 1128
flight potential,
age, 1
migration, 1
Pseudaletia unipuncta, 1
sex, 1
tethered-flight, 1
fluorescence,
chlorophyll, 215
raspberry, 215
Tetranychidae, 215
Tetranychus urticae, 215
food availability,
Aedes aegypti, 54
development, 54
mortality, 54
Plagiorchis elegans, 54
time-dependent transition probabilities, 54
food quality,
adult size, 168
Apanteles galleriae, 168
fecundity–sex ratio, 168
immature development, 168
longevity, 168
foodweb,
biocontrol, 388
host competition, 388
leafrollers, 388
shared parasitism, 388
Trigonospila brevifacies, 388
foraging, 47, 293, 977
Camponotus pennsylvanicus, 977
carpenter ant, 977
diet, 47
inbreeding, 293
liquid food, 47
nutrients, 977
social organization, 293
Solenopsis invicta, 47
solid food, 47
success rates, 47
territoriality, 293
forest,
caterpillar predation, 1097
invertebrate predators, 1097
larval dispersal, 1097
prey size, 1097
forest fire,
arthropod community, 774
disturbance, 774
Rio Grande, 774
riparian zone, 774
forest health,
Carabidae, 242
forest management, 242
Pinus ponderosa, 242
prescribed fire, 242
thinning, 242
forest litter arthropods,
Carabidae, 785
Oregon, 785
pitfall traps, 785
prescribed fire, 785
spiders, 785
forest management,
Carabidae, 242
forest health, 242
Pinus ponderosa, 242
prescribed fire, 242
thinning, 242
forest pest,
age-specific fecundity, 76
Anoplophora glabripennis, 76
exotic invasive species, 76
intrinsic growth rate, 76
Frankliniella fusca, 558, 1282
Allantonematidae, 558
biological control, 558
brachypterous, 1282
density dependence, 558
disease management, 1282
peanut, 558
population dynamics, 1282
Thripidae, 558
tospovirus, 1282
TSWV, 1282
Frankliniella occidentalis,
dendrobium orchids, 523
scouting methods, 523
thrips, 523

function,
community structure, 1052
leafminers, 1052
parasitoid assemblages, 1052
Phyllonorycter, 1052
Quercus, 1052
functional lower developmental thresholds,
Apthona nigriscutis, 348
biological control, 348
linear degree-day phenology models, 348
functional response,
Augmentative release, 1184
Chrysoperla rufilabris, 1184
Rhinocapsus vanduzeei, 1184
Stephanitis pyrioides, 1184
urban landscape, 1184
fungicidal effects,
biological control, 1206
entomopathogens, 1206
herbicide, 1206
nontarget effects, 1206
roundup, 1206

G

Galerucella calmariensis,
biological control, 880
Coleomegilla maculata, 880
larval behavior, 880
gene flow,
allozyme, 462
genetic structure, 462
Helicoverpa zea, 462
population genetics, 462
genetic structure,
allozyme, 462
gene flow, 462
Helicoverpa zea, 462
population genetics, 462
Geocoris punctipes,
Bacillus thuringiensis, 1197
biological control, 1197
Orius tristicolor, 1197
Spodoptera exigua, 1197
Zelus renardii, *1197*
geographical distribution,
CLIMEX, 1009
medfly, 1009
geostatistics, 149, 160, 273
Colorado potato beetle, 149
European corn borer, 160
global positioning systems, 149
insect spatial dynamics, 149
maize, 160
moisture, 273
Musca domestica, 273
population dynamics, 149
potatoes, 149
spatial autocorrelation, 273
spatial dynamics, 273
temporal dynamics, 273
glaciation,
Acari, 612
Azorella selago, 612
community structure, 612
habitat specificity, 612
Gleditsia japonica,
bruchid, 469
herbivorous insect, 469
overwintering stage, 469
seasonal adaptation, 469
voltinism, 469
global positioning systems,
Colorado potato beetle, 149
geostatistics, 149
insect spatial dynamics, 149
population dynamics, 149
potatoes, 149
glucobrassicin,
allylglucosinolate, 714
Brassicaceae, 714
induced defense, 714
specificity, 714
Glyptotendipes paripes, 608, 637
Chironomidae, 637
Cladotanytarsus, 637
development, 608
distribution, 637
life stages, 608
nuisance midge, 608
nuisance midges, 637
population biology, 637
temperature, 608
GNA,
biological control, 1219
Diatraea saccharalis, 1219
nontarget effects, 1219
snowdrop lectin, 1219
sugarcane, 1219
Goniozus,
Apanteles, 988
mode of parasitism, 988
temporal density-dependence, 988
tortricid leafrollers, 988
grape berry moth,
degree-day models, 708
diapause, 708
Endopiza viteana, 708
phenology, 708
grape phylloxera,
demography, 120
plant growth stage, 120
population growth, 120
Grapholita,
Acrobasis, 852
blueberry, 852
cherry fruitworm, 852
cranberry fruitworm, 852
Vaccinium, 852
grasshoppers,
Acrididae, 1158
barrier island, 1158
coastal vegetation, 1158
Orthoptera, 1158
gregarinidae,
biological control, 573
Istocheta, 573
Ovavesicula, 573
Popillia japonica, 573
Tiphia, 573
growth/differentiation balance,
fertilization, 923
host plant resistance, 923
secondary metabolism, 923
woody plants, 923
gypsy moth,
disparlure, 1119
Lymantria dispar, 1119
mating disruption, 1119
pheromone, 1119
release-recapture, 1119

H

habitat,
Camponotus vicinus, 1037
moisture, 1037
nest aggregation, 1037
nesting site, 1037
temperature, 1037
habitat specificity,
Acari, 612
Azorella selago, 612
community structure, 612
glaciation, 612
Halticoptera helioponi,
body size, 874
Chrysocharis flacilla, 874
Liriomyza huidobrensis, 874
Phaedrotoma scabriventris, 874
tritrophic interactions, 874
Hampea,
Anthonomus grandis, 693
dioecy, 693
Mexico, 693
reproductive ecology, 693
Harmonia axyridis,
biodiversity, 827
biological control, 827
competition, 827
intraguild predation, 827
multicolored Asian lady beetle, 827
Helicoverpa,
biomonitoring, 965
developmental instability, 965
environmental stress, 965
trait variability, 965
Helicoverpa zea, 232, 462
allozyme, 462
Coccinellidae, 232
diel activity, 232
diurnal, 232
gene flow, 462
genetic structure, 462

Hemiptera, 232
population genetics, 462
predation, 232
Hemiptera,
Coccinellidae, 232
diel activity, 232
diurnal, 232
Helicoverpa zea, 232
predation, 232
herbicide, 449, 1206
biological control, 1206
Collembola, 449
entomopathogens, 1206
fungicidal effects, 1206
microarthropods, 449
nontarget effects, 1206
population ecology, 449
roundup, 1206
soil mesofauna, 449
transgenic soybeans, 449
herbivore feeding,
leaf age, 836
leaf position, 836
Nicotiana tabacum, 836
nitrogen fertilization, 836
Populus deltoides, 836
herbivorous insect,
bruchid, 469
Gleditsia japonica, 469
overwintering stage, 469
seasonal adaptation, 469
voltinism, 469
herbivory,
behavioral ethogram, 538
electropenetration graph, 538
Lygus hesperus, 538
plant bugs, 538
probing behavior, 538
heritability,
bark beetle, 1276
directional selection, 1276
host acceptance, 1276
plant–insect interactions, 1276
Heterorhabditis bacteriophora, 895, 1240
alfalfa, 1240
clover root curculio, 1240
cranberry, 895
entomopathogenic nematodes, 1240
Heterorhabditis marelatus, 895
Otiorhynchus sulcatus, 895
Sitona hispidulus, 1240
Steinernema carpocapsae, 895
strawberry, 895
Heterorhabditis marelatus,
cranberry, 895
Heterorhabditis bacteriophora, 895
Otiorhynchus sulcatus, 895
Steinernema carpocapsae, 895
strawberry, 895
highbush blueberries,
Exomala orientalis, 1268
mating disruption, 1268
ornamental nurseries, 1268
sex pheromone, 1268
(Z)-7-tetradecen-2-one, 1268
Hippodamia convergens, 1168, 1191
Acheta domesticus, 1191
biological control, 1168
Chrysoperla carnea, 1168
Coccinella septempunctata, 1168
intraguild predation, 1168
Metarhizium anisopliae, 1191
nontarget effects, 1191
Oncopeltus fasciatus, 1191
pathogenicity, 1191
Solenopsis invicta, 1168
Homoeosoma electellum,
crop domestication, 626
native agricultural habitats, 626
parasitism, 626
sunflowers, 626
tritrophic interactions, 626
host,
biology, 354
demography, 354
parasitoid, 354
temperature, 354
host acceptance,
bark beetle, 1276
directional selection, 1276
heritability, 1276
plant–insect interactions, 1276
host age,
biological control, 505
host size, 505
sex ratio, 505
Spalangia cameroni, 505
host availability,
Cicadellidae, 914
desert plants, 914
leafhoppers, 914
host competition,
biocontrol, 388
foodweb, 388
leafrollers, 388
shared parasitism, 388
Trigonospila brevifacies, 388
host density,
feeding treatment, 732
muscoid flies, 732
parasitoid female age, 732
Tachinaephagus zealandicus, 732
temperature, 732
host discrimination,
alley cropping, 281
Conotrachelus retentus, 281
eastern black walnut, 281
Juglans nigra, 281
resource partitioning, 281
host effects,
Anoplophora glabripennis, 490
fecundity, 490
longevity, 490
host plant,
beet armyworm, 142
bertha armyworm, 1135
Mamestra configurata, 1135
oviposition, 1135
oviposition preference, 142
Spodoptera exigua, 142
host plant resistance, 22, 923
Empoasca fabae, 22
exudates, 22
fertilization, 923
growth/differentiation balance, 923
Medicago sativa, 22
plant volatiles, 22
secondary metabolism, 923
woody plants, 923
host size,
biological control, 505
host age, 505
sex ratio, 505
Spalangia cameroni, 505
hybrid fire ant,
black imported fire ant, 127
red imported fire ant, 127
Solenopsis invicta, 127
Solenopsis richteri, 127
Thelohania solenopsae, 127
Hylastes longicollis,
Dendroctonus ponderosae, 759
Hylurgops porosus, 759
Ips latidens, 759
Ips pini, 759
Pinus contorta, 759
Hylurgops porosus,
Dendroctonus ponderosae, 759
Hylastes longicollis, 759
Ips latidens, 759
Ips pini, 759
Pinus contorta, 759

I

Idiarthron subquadratum,
coffee pest, 516
mark-recapture, 516
population estimation, 516
sampling, 516
trapping, 516
immature development,
adult size, 168
Apanteles galleriae, 168
fecundity–sex ratio, 168
food quality, 168
longevity, 168

impact,
Cotesia glomerata, 367
Cotesia rubecula, 367
introduction, 367
Pieris rapae, 367
spread, 367
inbreeding,
foraging, 293
social organization, 293
territoriality, 293
indicator species,
biological assessment, 1142
Coleoptera, 1142
Orthoptera, 1142
induced defense,
allylglucosinolate, 714
Brassicaceae, 714
glucobrassicin, 714
specificity, 714
insect bioassay,
Bacillus thuringiensis, 30
Cry1Ac protein, 30
environmental risk, 30
enzyme-linked immunosorbent assays, 30
transgenic cotton, 30
insect spatial dynamics,
Colorado potato beetle, 149
geostatistics, 149
global positioning systems, 149
population dynamics, 149
potatoes, 149
insecticides,
biological control, 339
Dasineura oxycoccana, 339
parasitoids, 339
Syrphidae, 339
Vaccinium, 339
insect–plant interaction,
artificial diet, 306
Bemisia tabaci, 306
environmental parameters, 306
oviposition, 306
intensive forestry,
Aculops lobuliferus, 1000
defoliation, 1000
Populus deltoides, 1000
terminal mortality, 1000
interior spruce,
Mexican spruce, 198
Pissodes strobi, 198
Sitka spruce, 198
white pine weevil, 198
intracolonial prevalence,
biological control, 887
dual infections, 887
Solenopsis invicta, 887
Thelohania solenopsae, 887
Vairimorpha invictae, 887
intraguild predation, 827, 1168
biodiversity, 827
biological control, 827, 1168
Chrysoperla carnea, 1168
Coccinella septempunctata, 1168
competition, 827
Harmonia axyridis, 827
Hippodamia convergens, 1168
multicolored Asian lady beetle, 827
Solenopsis invicta, 1168
intrinsic growth rate,
age-specific fecundity, 76
Anoplophora glabripennis, 76
exotic invasive species, 76
forest pest, 76
introduction,
Cotesia glomerata, 367
Cotesia rubecula, 367
impact, 367
Pieris rapae, 367
spread, 367
invertebrate predators,
caterpillar predation, 1097
forest, 1097
larval dispersal, 1097
prey size, 1097
Ips,
conifer, 1043
Platysoma, 1043
population dynamics, 1043
predators, 1043
Thanasimus, 1043
Ips latidens,
Dendroctonus ponderosae, 759
Hylastes longicollis, 759
Hylurgops porosus, 759
Ips pini, 759
Pinus contorta, 759
Ips pini, 261, 759
Dendroctonus ponderosae, 759
exotic pest, 261
Hylastes longicollis, 759
Hylurgops porosus, 759
Ips latidens, 759
nonindigenous insect, 261
pheromone trapping, 261
Pinus contorta, 759
predators, 261
Thanasimus dubius, 261
Istocheta,
biological control, 573
gregarinidae, 573
Ovavesicula, 573
Popillia japonica, 573
Tiphia, 573
Ixodes scapularis,
Amblyomma americanum, 1085
flag sampling, 1085
Lyme disease, 1085

J

Juglans nigra,
alley cropping, 281
Conotrachelus retentus, 281
eastern black walnut, 281
host discrimination, 281
resource partitioning, 281

K

kairomone,
aggregation, 934
Cleridae, 766
energetic reserves, 941
Enoclerus sphegeus, 766
Macrodactylus subspinosus, 934
nutrition, 941
pharmacophagy, 941
pheromone, 934, 941
plant volatiles, 934
Scolytidae, 766
survival, 941
Thanasimus undatulus, 766
Trypodendron lineatum, 766
Kampimodromus aberrans,
domatia, 648
leaf trichomes, 648
pollen, 648
vineyards, 648

L

Lacanobia subjuncta,
apple, 995
degree-days, 995
development, 995
phenology model, 995
landscape structure,
alfalfa, 253
aphids, 253
Coccinellidae, 253
natural enemies, 253
predator-prey interaction, 253
larval behavior,
biological control, 880
Coleomegilla maculata, 880
Galerucella calmariensis, 880
larval dispersal,
caterpillar predation, 1097
forest, 1097
invertebrate predators, 1097
prey size, 1097
leaf age,
herbivore feeding, 836
leaf position, 836
Nicotiana tabacum, 836
nitrogen fertilization, 836
Populus deltoides, 836
leaf position,
herbivore feeding, 836
leaf age, 836
Nicotiana tabacum, 836
nitrogen fertilization, 836

Populus deltoides, 836
leaf trichomes,
domatia, 648
Kampimodromus aberrans, 648
pollen, 648
vineyards, 648
leafhoppers,
Cicadellidae, 914
desert plants, 914
host availability, 914
leafminers, 797, 1052
community structure, 1052
development, 797
function, 1052
Liriomyza huidobrensis, 797
Liriomyza trifolii, 797
mortality, 797
parasitoid assemblages, 1052
Phyllonorycter, 1052
Quercus, 1052
temperature dependence, 797
leafrollers,
biocontrol, 388
foodweb, 388
host competition, 388
shared parasitism, 388
Trigonospila brevifacies, 388
Leiobunum,
alfalfa, 92
Phalangium opilio, 92
phenology, 92
population, 92
soybean, 92
Leptinotarsa decemlineata,
density, 533
dispersal, 533
flight, 533
Leptoglossus occidentalis,
abortion, 1023
conelets, 1023
Pinus contorta var. *latifolia*, 1023
Pinus monticola, 1023
seed loss, 1023
life history,
oviposition, 702
photoperiod, 702
reproductive output, 702
season, 702
time constraints, 702
life stages,
development, 608
Glyptotendipes paripes, 608
nuisance midge, 608
temperature, 608
life table,
biological control, 482
Colorado potato beetle, 1110
Ostrinia nubilalis, 482
population dynamics, 1110
potatoes, 1110
tomatoes, 1110
Trichogramma, 482
linear degree-day phenology models,
Apthona nigriscutis, 348
biological control, 348
functional lower developmental thresholds, 348
liquid food,
diet, 47
foraging, 47
Solenopsis invicta, 47
solid food, 47
success rates, 47
Liriomyza huidobrensis, 797, 874
body size, 874
Chrysocharis flacilla, 874
development, 797
Halticoptera helioponi, 874
leafminers, 797
Liriomyza trifolii, 797
mortality, 797
Phaedrotoma scabriventris, 874
temperature dependence, 797
tritrophic interactions, 874
Liriomyza trifolii,
development, 797
leafminers, 797
Liriomyza huidobrensis, 797
mortality, 797
temperature dependence, 797
Listronotus oregonensis,
decision rules, 1251
Petroselinum crispum, 1251
probability density functions, 1251
spatial pattern, 1251
transect pattern, 1251
longevity, 168, 375, 432, 490
adult size, 168
Anoplophora glabripennis, 490
Ayanteles galleriae, 168
biological control, 432
development rate, 375, 432
Eoreuma loftini, 432
fecundity, 490
fecundity–sex ratio, 168
food quality, 168
host effects, 490
immature development, 168
Lydella jalisco, 432
muscoid flies, 375
stalkborer, 432
Tachinaephagus zealandicus, 375
temperature, 375
lower developmental thresholds,
adult longevity, 804
Cydia pomonella, 804
degree-days, 804
fecundity, 804
oviposition behavior, 804
Lydella jalisco,
biological control, 432
development rate, 432
Eoreuma loftini, 432
longevity, 432
stalkborer, 432
Lygus hesperus,
behavioral ethogram, 538
electropenetration graph, 538
herbivory, 538
plant bugs, 538
probing behavior, 538
Lymantria dispar, 668, 1071, 1119
computer model, 1071
disparlure, 1119
gypsy moth, 1119
mast, 668
mating disruption, 1119
pathogen, 1071
Peromyscus, 668
pheromone, 1119
predation, 668
release-recapture, 1119
resting spores, 1071
Lyme disease,
Amblyomma americanum, 1085
flag sampling, 1085
Ixodes scapularis, 1085
Lysiphlebia mirzai,
biological control, 418
brown citrus aphid, 418
temperature-dependent development, 418
Lysiphlebus testaceipes,
azadirachtin, 172
brown citrus aphid, 172
neem extract, 172
Neemix, 172
Toxoptera citricida, 172

M

Macrodactylus subspinosus,
aggregation, 934
kairomone, 934
pheromone, 934
plant volatiles, 934
Macrolophus pygmaeus,
developmental temperature threshold, 661
egg development, 661
Miridae, 661
nymphal development, 661
thermal constant, 661
maize,
European corn borer, 160
geostatistics, 160
Mamestra configurata,
bertha armyworm, 1135
host plant, 1135
oviposition, 1135

manure,
development, 588
Diptera, 588
Fannia, 588
moisture, 588
mark recapture, 65, 516
biological control, 425
Brassica spp., 65
coffee pest, 516
dispersal, 425
Idiarthron subquadratum, 516
movement, 65
Musca domestica, 425
Phthorimaea operculella, 65
Plutella xylostella, 65
population estimation, 516
potatoes, 65
sampling, 516
Spalangia cameroni, 425
spatial distribution, 425
trapping, 516
mass trapping,
China, 602
Cydia trasias, 602
sex pheromone, 602
Sophora japonica, 602
tortrix, 602
mast,
Lymantria dispar, 668
Peromyscus, 668
predation, 668
mating disruption, 1119, 1268
disparlure, 1119
Exomala orientalis, 1268
gypsy moth, 1119
highbush blueberries, 1268
Lymantria dispar, 1119
ornamental nurseries, 1268
pheromone, 1119
release-recapture, 1119
sex pheromone, 1268
(Z)-7-tetradecen-2-one, 1268
medfly,
CLIMEX, 1009
geographical distribution, 1009
Medicago sativa,
Empoasca fabae, 22
exudates, 22
host plant resistance, 22
plant volatiles, 22
Metarhizium anisopliae,
Acheta domesticus, 1191
Hippodamia convergens, 1191
nontarget effects, 1191
Oncopeltus fasciatus, 1191
pathogenicity, 1191
Mexican spruce,
interior spruce, 198
Pissodes strobi, 198
Sitka spruce, 198
white pine weevil, 198
Mexico,
Anthonomus grandis, 693
dioecy, 693
Hampea, 693
reproductive ecology, 693
microarthropods,
Collembola, 449
herbicide, 449
population ecology, 449
soil mesofauna, 449
transgenic soybeans, 449
microbial control,
Beauveria bassiana, 746
entomopathogenic fungi, 746
red imported fire ant, 746
Solenopsis invicta, 746
microclimate,
augmentative biological control, 361
cotton, 361
microhabitat, 361
Trichogramma augmentation, 361
microhabitat,
augmentative biological control, 361
cotton, 361
microclimate, 361
Trichogramma augmentation, 361
microsporidia,
biological control, 1091
colony size, 1091
fire ants, 1091
Thelohania solenopsae, 1091
migration,
age, 1
flight potential, 1
Pseudaletia unipuncta, 1
sex, 1
tethered-flight, 1
Miridae,
developmental temperature threshold, 661
egg development, 661
Macrolophus pygmaeus, 661
nymphal development, 661
thermal constant, 661
mode of parasitism,
Apanteles, 988
Goniozus, 988
temporal density-dependence, 988
tortricid leafrollers, 988
model,
Acrobasis nuxvorella, 1062
Bacillus thuringiensis, 208
degradation, 208
pecan, 1062
Pecan nut casebearer, 1062
soil, 208
strain PS149B1, 208
moisture, 273, 588, 1037
Camponotus vicinus, 1037
development, 588
Diptera, 588
Fannia, 588
geostatistics, 273
habitat, 1037
manure, 588
Musca domestica, 273
nest aggregation, 1037
nesting site, 1037
spatial autocorrelation, 273
spatial dynamics, 273
temperature, 1037
temporal dynamics, 273
mole crickets,
behavior, 476
damage, 476
Scapteriscus, 476
soil moisture, 476
surface activity, 476
montane forest spiders,
sampling intensity, 319
sampling methods, 319
species richness estimation, 319
mortality, 54, 797, 816
Aedes aegypti, 54
Cryptolestes ferrugineus, 816
development, 54, 797
food availability, 54
leafminers, 797
Liriomyza huidobrensis, 797
Liriomyza trifolii, 797
movement, 816
Plagiorchis elegans, 54
stored-grain, 816
temperature dependence, 797
temperature gradient, 816
time-dependent transition probabilities, 54
mosquito larvae,
Anopheles, 134
breeding, 134
vegetation, 134
water velocity, 134
movement, 65, 816
Brassica spp., 65
Cryptolestes ferrugineus, 816
mark-recapture, 65
mortality, 816
Phthorimaea operculella, 65
Plutella xylostella, 65
potatoes, 65
stored-grain, 816
temperature gradient, 816
multicolored Asian lady beetle,
biodiversity, 827
biological control, 827
competition, 827
Harmonia axyridis, 827
intraguild predation, 827

multiparasitism,
entomopathogenic nematode, 1226
Ormia depleta, 1226
Scapteriscus spp., 1226
Steinernema scapterisci, 1226
tachinid, 1226
Musca domestica, 273, 411, 425
biological control, 425
Dihrinus himalayanus, 411
dispersal, 425
geostatistics, 273
mark recapture, 425
moisture, 273
Muscidifurax raptor, 411
Spalangia cameroni, 411, 425
Spalangia endius, 411
Spalangia gemina, 411
spatial autocorrelation, 273
spatial distribution, 425
spatial dynamics, 273
temporal dynamics, 273
Muscidifurax raptor,
Dihrinus himalayanus, 411
Musca domestica, 411
Spalangia cameroni, 411
Spalangia endius, 411
Spalangia gemina, 411
muscoid flies, 375, 732
development rate, 375
feeding treatment, 732
host density, 732
longevity, 375
parasitoid female age, 732
Tachinaephagus zealandicus, 375, 732
temperature, 375, 732

N

native agricultural habitats,
crop domestication, 626
Homoeosoma electellum, 626
parasitism, 626
sunflowers, 626
tritrophic interactions, 626
natural enemies,
alfalfa, 253
aphids, 253
Coccinellidae, 253
landscape structure, 253
predator-prey interaction, 253
natural mortality,
Anthonomus grandis grandis, 972
boll weevil, 972
critical weight, 972
neem extract,
azadirachtin, 172
brown citrus aphid, 172
Lysiphlebus testaceipes, 172
Neemix, 172
Toxoptera citricida, 172
Neemix,
azadirachtin, 172
brown citrus aphid, 172
Lysiphlebus testaceipes, 172
neem extract, 172
Toxoptera citricida, 172
nest aggregation,
Camponotus vicinus, 1037
habitat, 1037
moisture, 1037
nesting site, 1037
temperature, 1037
nesting site,
Camponotus vicinus, 1037
habitat, 1037
moisture, 1037
nest aggregation, 1037
temperature, 1037
Nicotiana tabacum,
herbivore feeding, 836
leaf age, 836
leaf position, 836
nitrogen fertilization, 836
Populus deltoides, 836
nitrogen fertilization,
herbivore feeding, 836
leaf age, 836
leaf position, 836
Nicotiana tabacum, 836
Populus deltoides, 836
nitrogen limitation,
fecundity, 313
ovaries, 313
reproduction, 313
termites, 313
Zootermopsis angusticollis, 313
nonindigenous insect,
exotic pest, 261
Ips pini, 261
pheromone trapping, 261
predators, 261
Thanasimus dubius, 261
nonlinear models,
developmental rate, 177
predator-prey synchrony, 177
Stethorus punctillum, 177
temperature, 177
Tetranychus mcdanieli, 177
nontarget effects, 1191, 1206, 1219
Acheta domesticus, 1191
biological control, 1206, 1219
Diatraea saccharalis, 1219
entomopathogens, 1206
fungicidal effects, 1206
GNA, 1219
herbicide, 1206
Hippodamia convergens, 1191
Metarhizium anisopliae, 1191
Oncopeltus fasciatus, 1191
pathogenicity, 1191
roundup, 1206
snowdrop lectin, 1219
sugarcane, 1219
nontarget impacts,
California, 723
Cydia pomonella, 723
parasitoid, 723
walnuts, 723
nontarget Lepidoptera,
Bacillus thuringiensis subsp. *kurstaki*, 903
Choristoneura occidentalis, 903
Douglas-fir, 903
western spruce budworm, 903
nuisance midge,
Chironomidae, 637
Cladotanytarsus, 637
development, 608
distribution, 637
Glyptotendipes paripes, 608, 637
life stages, 608
population biology, 637
temperature, 608
nutrients,
Camponotus pennsylvanicus, 977
carpenter ant, 977
foraging, 977
nutrition, 99, 941
caste, 99
energetic reserves, 941
kairomones, 941
parental manipulation, 99
pharmacophagy, 941
phenotypic plasticity, 99
pheromones, 941
Polistes metricus, 99
social wasp, 99
survival, 941
nymphal development,
developmental temperature threshold, 661
egg development, 661
Macrolophus pygmaeus, 661
Miridae, 661
thermal constant, 661

O

obliquebanded leafroller,
codling moth, 960
oriental fruit moth, 960
pheromone, 960
piezoelectric sprayer, 960
tortricids, 960
Oncopeltus fasciatus,
Acheta domesticus, 1191
Hippodamia convergens, 1191
Metarhizium anisopliae, 1191
nontarget effects, 1191
pathogenicity, 1191

orchard,
attracticide, 189
behavior, 189
Cydia pomonella, 189
pheromone, 189
Oregon,
Carabidae, 785
forest litter arthropods, 785
pitfall traps, 785
prescribed fire, 785
spiders, 785
Orgyia leucostigma,
Betula papyrifera, 10
condensed tannin, 10
paper birch, 10
whitemarked tussock moth, 10
oriental fruit moth,
codling moth, 960
obliquebanded leafroller, 960
pheromone, 960
piezoelectric sprayer, 960
tortricids, 960
Orius tristicolor,
Bacillus thuringiensis, 1197
biological control, 1197
Geocoris punctipes, 1197
Spodoptera exigua, 1197
Zelus renardii, 1197
Ormia depleta,
entomopathogenic nematode, 1226
multiparasitism, 1226
Scapteriscus spp., 1226
Steinernema scapterisci, 1226
tachinid, 1226
ornamental nurseries,
Exomala orientalis, 1268
highbush blueberries, 1268
mating disruption, 1268
sex pheromone, 1268
(Z)-7-tetradecen-2-one, 1268
ornamentals,
conservation biological control, 564
parasitoids, 564
predators, 564
turfgrass, 564
wildflowers, 564
Orthoptera, 1142, 1158
Acrididae, 1158
barrier island, 1158
biological assessment, 1142
coastal vegetation, 1158
Coleoptera, 1142
grasshoppers, 1158
indicator species, 1142
Ostrinia nubilalis, 482, 676
biological control, 482
Diatraea grandiosella, 676
European corn borer, 676
life tables, 482
resistance management, 676
simulation modeling, 676
southwestern corn borer, 676
Trichogramma, 482
Otiorhynchus sulcatus,
cranberry, 895
Heterorhabditis bacteriophora, 895
Heterorhabditis marelatus, 895
Steinernema carpocapsae, 895
strawberry, 895
ovaries,
fecundity, 313
nitrogen limitation, 313
reproduction, 313
termites, 313
Zootermopsis angusticollis, 313
Ovavesicula,
biological control, 573
gregarinidae, 573
Istocheta, 573
Popillia japonica, 573
Tiphia, 573
overwintering stage,
bruchid, 469
Gleditsia japonica, 469
herbivorous insect, 469
seasonal adaptation, 469
voltinism, 469
oviposition, 107, 112, 285, 306, 702, 1135
Abutilon theophrasti, 107, 112
Acalypha ostryaefolia, 107, 112
artificial diet, 306
Bemisia tabaci, 306
bertha armyworm, 1135
Coleomegilla maculata, 107, 112
crop rotation, 285
environmental parameters, 306
host plant, 1135
insect–plant interaction, 306
life history, 702
Mamestra configurata, 1135
photoperiod, 702
predation, 112
reproductive output, 702
resistance, 285
season, 702
time constraints, 702
whitefly, 107
Zea mays, 107, 112
oviposition behavior, 804, 859
Acanthoscelides obtectus, 859
adult longevity, 804
beans, 859
Bruchidae, 859
Cydia pomonella, 804
degree-days, 804
fecundity, 804
lower developmental thresholds, 804
parasitoids, 859
storage pest, 859
oviposition preference,
beet armyworm, 142
host plants, 142
Spodoptera exigua, 142

P

paper birch,
Betula papyrifera, 10
condensed tannin, 10
Orgyia leucostigma, 10
whitemarked tussock moth, 10
parasite acclimation,
Bathyplectes curculionis, 739
cold tolerance, 739
parasitism,
crop domestication, 626
Homoeosoma electellum, 626
native agricultural habitats, 626
sunflowers, 626
tritrophic interactions, 626
parasitoid, 339, 354, 397, 403, 564, 723, 859, 947
Acanthoscelides obtectus, 859
Argentina, 403
beans, 859
biological control, 339, 397, 403, 947
biology, 354
Bruchidae, 859
California, 723
codling moth, 947
conservation biological control, 564
Cydia pomonella, 723, 947
Dasineura oxycoccana, 339
demography, 354
developmental times, 403
Eretmocerus, 397
fire ant, 403
host, 354
insecticides, 339
nontarget impacts, 723
ornamentals, 564
oviposition behavior, 859
phenology, 947
pheromone, 947
predators, 564
storage pest, 859
Syrphidae, 339
temperature, 354
temperature effects, 403
tomato, 397
tritrophic, 397
turfgrass, 564
Vaccinium, 339
walnuts, 723
whitefly, 397
wildflowers, 564

parasitoid assemblages,
community structure, 1052
function, 1052
leafminers, 1052
Phyllonorycter, 1052
Quercus, 1052
parasitoid female age,
feeding treatment, 732
host density, 732
muscoid flies, 732
Tachinaephagus zealandicus, 732
temperature, 732
parental manipulation,
caste, 99
nutrition, 99
phenotypic plasticity, 99
Polistes metricus, 99
social wasp, 99
pathogen,
computer model, 1071
Lymantria dispar, 1071
resting spores, 1071
pathogenicity,
Acheta domesticus, 1191
Hippodamia convergens, 1191
Metarhizium anisopliae, 1191
nontarget effects, 1191
Oncopeltus fasciatus, 1191
peach,
apple, 1259
Conotrachelus, 1259
plum curculio, 1259
sampling, 1259
trap, 1259
peanut,
Allantonematidae, 558
biological control, 558
density dependence, 558
Frankliniella fusca, 558
Thripidae, 558
pecan,
Acrobasis nuxvorella, 1062
models, 1062
Pecan nut casebearer, 1062
Pecan nut casebearer,
Acrobasis nuxvorella, 1062
models, 1062
pecan, 1062
Peromyscus,
Lymantria dispar, 668
mast, 668
predation, 668
Petroselinum crispum,
decision rules, 1251
Listronotus oregonensis, 1251
probability density functions, 1251
spatial pattern, 1251
transect pattern, 1251
Phaedrotoma scabriventris,
body size, 874
Chrysocharis flacilla, 874
Halticoptera helioponi, 874
Liriomyza huidobrensis, 874
tritrophic interactions, 874
Phalangium opilio,
alfalfa, 92
Leiobunum, 92
phenology, 92
population, 92
soybean, 92
pharmacophagy,
energetic reserves, 941
kairomones, 941
nutrition, 941
pheromones, 941
survival, 941
phenology, 92, 708, 947
alfalfa, 92
biological control, 947
codling moth, 947
Cydia pomonella, 947
degree-day models, 708
diapause, 708
Endopiza viteana, 708
grape berry moth, 708
Leiobunum, 92
parasitoid, 947
Phalangium opilio, 92
pheromone, 947
population, 92
soybean, 92
phenology model,
apple, 995
degree-days, 995
development, 995
Lacanobia subjuncta, 995
phenotypic plasticity,
caste, 99
nutrition, 99
parental manipulation, 99
Polistes metricus, 99
social wasp, 99
pheromone, 189, 934, 947, 960, 1119
aggregation, 934
attracticide, 189
behavior, 189
biological control, 947
codling moth, 947, 960
Cydia pomonella, 189, 947
disparlure, 1119
energetic reserves, 941
gypsy moth, 1119
kairomone, 934, 941
Lymantria dispar, 1119
Macrodactylus subspinosus, 934
mating disruption, 1119
nutrition, 941
obliquebanded leafroller, 960
orchard, 189
oriental fruit moth, 960
parasitoid, 947
pharmacophagy, 941
phenology, 947
piezoelectric sprayer, 960
plant volatiles, 934
release-recapture, 1119
survival, 941
tortricids, 960
pheromone trapping,
exotic pest, 261
Ips pini, 261
nonindigenous insect, 261
predators, 261
Thanasimus dubius, 261
pheromone traps,
Anthonomus grandis, 1128
boll weevil, 1128
flight behavior, 1128
sampling, 1128
wind, 1128
photoperiod,
life history, 702
oviposition, 702
reproductive output, 702
season, 702
time constraints, 702
Phthorimaea operculella,
Brassica spp., 65
mark-recapture, 65
movement, 65
Plutella xylostella, 65
potatoes, 65
Phyllonorycter,
community structure, 1052
function, 1052
leafminers, 1052
parasitoid assemblages, 1052
Quercus, 1052
phytoextraction,
alfalfa, 953
biomagnification, 953
phytoremediation, 953
selenium, 953
Spodoptera exigua, 953
phytoremediation,
alfalfa, 953
biomagnification, 953
phytoextraction, 953
selenium, 953
Spodoptera exigua, 953
Pieris rapae, 367, 1231
Bacillus thuringiensis, 1231
biological control, 1231
cabbage, 1231
Cotesia glomerata, 367
Cotesia rubecula, 367
impact, 367
introduction, 367
spread, 367
sugar sprays, 1231

piezoelectric sprayer,
codling moth, 960
obliquebanded leafroller, 960
oriental fruit moth, 960
pheromone, 960
tortricids, 960
Pinus contorta,
Dendroctonus ponderosae, 759
Hylastes longicollis, 759
Hylurgops porosus, 759
Ips latidens, 759
Ips pini, 759
Pinus contorta var. *latifolia*,
abortion, 1023
conelets, 1023
Leptoglossus occidentalis, 1023
Pinus monticola, 1023
seed loss, 1023
Pinus monticola,
abortion, 1023
conelets, 1023
Leptoglossus occidentalis, 1023
Pinus contorta var. *latifolia*, 1023
seed loss, 1023
Pinus ponderosa,
Carabidae, 242
forest health, 242
forest management, 242
prescribed fire, 242
thinning, 242
Pissodes strobi,
interior spruce, 198
Mexican spruce, 198
Sitka spruce, 198
white pine weevil, 198
pitfall trap size,
Araneae, 438
biodiversity, 438
Carabidae, 438
sampling, 438
Staphylindae, 438
pitfall traps,
Carabidae, 785
forest litter arthropods, 785
Oregon, 785
prescribed fire, 785
spiders, 785
Plagiorchis elegans,
Aedes aegypti, 54
development, 54
food availability, 54
mortality, 54
time-dependent transition probabilities, 54
plant bugs,
behavioral ethogram, 538
electropenetration graph, 538
herbivory, 538
Lygus hesperus, 538
probing behavior, 538
plant growth stage,
demography, 120
grape phylloxera, 120
population growth, 120
Plant structure,
connectivity, 1153
size, 1153
Trichogramma evanescens, 1153
trophic interactions, 1153
plant volatiles, 22, 934
aggregation, 934
Empoasca fabae, 22
exudates, 22
host plant resistance, 22
kairomone, 934
Macrodactylus subspinosus, 934
Medicago sativa, 22
pheromone, 934
plant–insect interactions,
bark beetle, 1276
directional selection, 1276
heritability, 1276
host acceptance, 1276
Platysoma,
conifer, 1043
Ips, 1043
population dynamics, 1043
predators, 1043
Thanasimus, 1043
plum curculio,
apple, 1259
Conotrachelus, 1259
peach, 1259
sampling, 1259
trap, 1259
Plutella xylostella, 65, 221
Brassica spp., 65
development time, 221
mark-recapture, 65
movement, 65
Phthorimaea operculella, 65
potatoes, 65
rate function, 221
survival, 221
temperature, 221
Polistes metricus,
caste, 99
nutrition, 99
parental manipulation, 99
phenotypic plasticity, 99
social wasp, 99
pollen,
domatia, 648
Kampimodromus aberrans, 648
leaf trichomes, 648
vineyards, 648
Popillia japonica,
biological control, 573
gregarinidae, 573
Istocheta, 573
Ovavesicula, 573
Tiphia, 573
population,
alfalfa, 92
Leiobunum, 92
Phalangium opilio, 92
phenology, 92
soybean, 92
population biology,
Chironomidae, 637
Cladotanytarsus, 637
distribution, 637
Glyptotendipes paripes, 637
nuisance midges, 637
population dynamics, 149, 299, 686, 1043, 1110, 1282
brachypterous, 1282
Carposina sasakii, 686
climatic factors, 299
Colorado potato beetle, 149, 1110
conifer, 1043
Deois flavopicta, 299
disease management, 1282
Frankliniella fusca, 1282
geostatistics, 149
global positioning systems, 149
insect spatial dynamics, 149
Ips, 1043
life table, 1110
Platysoma, 1043
potatoes, 149, 1110
predators, 1043
seasonality, 299
survivorship, 299, 686
Thanasimus, 1043
tomatoes, 1110
tospovirus, 1282
TSWV, 1282
population ecology,
Collembola, 449
herbicide, 449
microarthropods, 449
soil mesofauna, 449
transgenic soybeans, 449
population estimation,
coffee pest, 516
Idiarthron subquadratum, 516
mark-recapture, 516
sampling, 516
trapping, 516
population genetics,
allozyme, 462
gene flow, 462
genetic structure, 462
Helicoverpa zea, 462
population growth,
demography, 120
grape phylloxera, 120
plant growth stage, 120
Populus deltoides, 836, 1000
Aculops lobuliferus, 1000
defoliation, 1000
herbivore feeding, 836

intensive forestry, 1000
leaf age, 836
leaf position, 836
Nicotiana tabacum, 836
nitrogen fertilization, 836
terminal mortality, 1000
postdiapause,
blueberry maggot, 1103
development rates, 1103
diapause, 1103
Rhagoletis mendax, 1103
potatoes, 65, 149, 1110
Brassica spp., 65
Colorado potato beetle, 149, 1110
geostatistics, 149
global positioning systems, 149
insect spatial dynamics, 149
life table, 1110
mark-recapture, 65
movement, 65
Phthorimaea operculella, 65
Plutella xylostella, 65
population dynamics, 149, 1110
tomatoes, 1110
predation, 112, 232, 668
Abutilon theophrasti, 112
Acalypha ostryaefolia, 112
Coccinellidae, 232
Coleomegilla maculata, 112
diel activity, 232
diurnal, 232
Helicoverpa zea, 232
Hemiptera, 232
Lymantria dispar, 668
mast, 668
oviposition, 112
Peromyscus, 668
Zea mays, 112
predator interference,
ant-aphid mutualism, 1175
Aphis gossypii, 1175
Chrysoperla carnea, 1175
Coccinellidae, 1175
Solenopsis invicta, 1175
predator-prey interaction,
alfalfa, 253
aphids, 253
Coccinellidae, 253
landscape structure, 253
natural enemies, 253
predator-prey synchrony,
developmental rate, 177
nonlinear models, 177
Stethorus punctillum, 177
temperature, 177
Tetranychus mcdanieli, 177
predators, 261, 564, 1043
conifer, 1043
conservation biological control, 564
exotic pest, 261
Ips, 1043
Ips pini, 261
nonindigenous insect, 261
ornamentals, 564
parasitoids, 564
pheromone trapping, 261
Platysoma, 1043
population dynamics, 1043
Thanasimus, 1043
Thanasimus dubius, 261
turfgrass, 564
wildflowers, 564
prescribed fire, 242, 785
Carabidae, 242, 785
forest health, 242
forest litter arthropods, 785
forest management, 242
Oregon, 785
Pinus ponderosa, 242
pitfall traps, 785
spiders, 785
thinning, 242
prey size,
caterpillar predation, 1097
forest, 1097
invertebrate predators, 1097
larval dispersal, 1097
probability density functions,
decision rules, 1251
Listronotus oregonensis, 1251
Petroselinum crispum, 1251
spatial pattern, 1251
transect pattern, 1251
probing behavior,
behavioral ethogram, 538
electropenetration graph, 538
herbivory, 538
Lygus hesperus, 538
plant bugs, 538
Pseudaletia unipuncta,
age, 1
flight potential, 1
migration, 1
sex, 1
tethered-flight, 1

Q

Quercus,
community structure, 1052
function, 1052
leafminers, 1052
parasitoid assemblages, 1052
Phyllonorycter, 1052

R

raspberry,
chlorophyll, 215
fluorescence, 215
Tetranychidae, 215
Tetranychus urticae, 215
rate function,
development time, 221
Plutella xylostella, 221
survival, 221
temperature, 221
red imported fire ant, 127, 746
Beauveria bassiana, 746
black imported fire ant, 127
entomopathogenic fungi, 746
hybrid fire ant, 127
microbial control, 746
Solenopsis invicta, 127, 746
Solenopsis richteri, 127
Thelohania solenopsae, 127
red sunflower seed weevil,
biological control, 844
Braconidae, 844
Curculionidae, 844
Smicronyx fulvus, 844
Triaspis aequoris, 844
release-recapture,
disparlure, 1119
gypsy moth, 1119
Lymantria dispar, 1119
mating disruption, 1119
pheromone, 1119
reproduction,
fecundity, 313
nitrogen limitation, 313
ovaries, 313
termites, 313
Zootermopsis angusticollis, 313
reproductive ecology,
Anthonomus grandis, 693
dioecy, 693
Hampea, 693
Mexico, 693
reproductive output,
life history, 702
oviposition, 702
photoperiod, 702
season, 702
time constraints, 702
resistance,
crop rotation, 285
oviposition, 285
resistance management,
Diatraea grandiosella, 676
European corn borer, 676
Ostrinia nubilalis, 676
simulation modeling, 676
southwestern corn borer, 676
resource partitioning,
alley cropping, 281
Conotrachelus retentus, 281
eastern black walnut, 281
host discrimination, 281
Juglans nigra, 281
resting spores,
computer model, 1071
Lymantria dispar, 1071
pathogen, 1071

Reticulitermes flavipes,
biological control, 381
Coptotermes formosanus, 381
entomopathogenic nematodes, 381
Rhagoletis mendax,
blueberry maggot, 1103
development rates, 1103
diapause, 1103
postdiapause, 1103
Rhinocapsus vanduzeei,
Augmentative release, 1184
Chrysoperla rufilabris, 1184
functional response, 1184
Stephanitis pyrioides, 1184
urban landscape, 1184
Rio Grande,
arthropod community, 774
disturbance, 774
forest fire, 774
riparian zone, 774
riparian zone,
arthropod community, 774
disturbance, 774
forest fire, 774
Rio Grande, 774
roundup,
biological control, 1206
entomopathogens, 1206
fungicidal effects, 1206
herbicide, 1206
nontarget effects, 1206

S

sampling, 438, 516, 1128, 1259
Anthonomus grandis, 1128
apple, 1259
Araneae, 438
biodiversity, 438
boll weevil, 1128
Carabidae, 438
coffee pest, 516
Conotrachelus, 1259
flight behavior, 1128
Idiarthron subquadratum, 516
mark-recapture, 516
peach, 1259
pheromone traps, 1128
pitfall trap size, 438
plum curculio, 1259
population estimation, 516
Staphylindae, 438
trap, 1259
trapping, 516
wind, 1128
sampling intensity,
montane forest spiders, 319
sampling methods, 319
species richness estimation, 319
sampling methods,
montane forest spiders, 319
sampling intensity, 319
species richness estimation, 319
Scapteriscus,
behavior, 476
damage, 476
mole crickets, 476
soil moisture, 476
surface activity, 476
Scapteriscus spp.,
entomopathogenic nematode, 1226
multiparasitism, 1226
Ormia depleta, 1226
Steinernema scapterisci, 1226
tachinid, 1226
Scolytidae,
Cleridae, 766
Enoclerus sphegeus, 766
Kairomone, 766
Thanasimus undatulus, 766
Trypodendron lineatum, 766
scouting methods,
dendrobium orchids, 523
Frankliniella occidentalis, 523
thrips, 523
season,
life history, 702
oviposition, 702
photoperiod, 702
reproductive output, 702
time constraints, 702
seasonal adaptation,
bruchid, 469
Gleditsia japonica, 469
herbivorous insect, 469
overwintering stage, 469
voltinism, 469
seasonality,
climatic factors, 299
Deois flavopicta, 299
population dynamics, 299
survivorship, 299
secondary metabolism,
fertilization, 923
growth/differentiation balance, 923
host plant resistance, 923
woody plants, 923
seed loss,
abortion, 1023
conelets, 1023
Leptoglossus occidentalis, 1023
Pinus contorta var. *latifolia*, 1023
Pinus monticola, 1023
selenium,
alfalfa, 953
biomagnification, 953
phytoextraction, 953
phytoremediation, 953
Spodoptera exigua, 953
semiochemicals,
Arachis hypogaea, 15
attractants, 15
carbon dioxide, 15
Elasmopalpus lignosellus, 15
sex,
age, 1
flight potential, 1
migration, 1
Pseudaletia unipuncta, 1
tethered-flight, 1
sex pheromone, 602, 1268
China, 602
Cydia trasias, 602
Exomala orientalis, 1268
highbush blueberries, 1268
mass trapping, 602
mating disruption, 1268
ornamental nurseries, 1268
Sophora japonica, 602
(*Z*)-7-tetradecen-2-one, 1268
tortrix, 602
sex ratio,
biological control, 505
host age, 505
host size, 505
Spalangia cameroni, 505
shared parasitism,
biocontrol, 388
foodweb, 388
host competition, 388
leafrollers, 388
Trigonospila brevifacies, 388
simulation modeling,
Diatraea grandiosella, 676
European corn borer, 676
Ostrinia nubilalis, 676
resistance management, 676
southwestern corn borer, 676
Sitka spruce,
interior spruce, 198
Mexican spruce, 198
Pissodes strobi, 198
white pine weevil, 198
Sitona hispidulus,
alfalfa, 1240
clover root curculio, 1240
entomopathogenic nematodes, 1240
Heterorhabditis bacteriophora, 1240
size,
connectivity, 1153
Plant structure, 1153
Trichogramma evanescens, 1153
trophic interactions, 1153
Smicronyx fulvus,
biological control, 844
Braconidae, 844
Curculionidae, 844

red sunflower seed weevil, 844
Triaspis aequoris, 844
snowdrop lectin,
biological control, 1219
Diatraea saccharalis, 1219
GNA, 1219
nontarget effects, 1219
sugarcane, 1219
social organization,
foraging, 293
inbreeding, 293
territoriality, 293
social wasp,
caste, 99
nutrition, 99
parental manipulation, 99
phenotypic plasticity, 99
Polistes metricus, 99
soil,
Bacillus thuringiensis, 208
degradation, 208
model, 208
strain PS149B1, 208
soil ecology,
agricultural disturbance, 37
Collembola, 37
cropping systems, 37
springtails, 37
soil mesofauna,
Collembola, 449
herbicide, 449
microarthropods, 449
population ecology, 449
transgenic soybeans, 449
soil moisture,
behavior, 476
damage, 476
mole crickets, 476
Scapteriscus, 476
surface activity, 476
soldier proportion,
Coptotermes formosanus, 509
temperature, 509
termite survival, 509
wood consumption, 509
Solenopsis invicta, 47, 127, 746, 887, 1168, 1175
ant-aphid mutualism, 1173
Aphis gossypii, 1175
Beauveria bassiana, 746
biological control, 887, 1168
black imported fire ant, 127
Chrysoperla carnea, 1168, 1175
Coccinella septempunctata, 1168
Coccinellidae, 1175
diet, 47
dual infections, 887
entomopathogenic fungi, 746
foraging, 47
Hippodamia convergens, 1168
hybrid fire ant, 127
intracolonial prevalence, 887
intraguild predation, 1168
liquid food, 47
microbial control, 746
predator interference, 1175
red imported fire ant, 127, 746
Solenopsis richteri, 127
solid food, 47
success rates, 47
Thelohania solenopsae, 127, 887
Vairimorpha invictae, 887
Solenopsis richteri,
black imported fire ant, 127
hybrid fire ant, 127
red imported fire ant, 127
Solenopsis invicta, 127
Thelohania solenopsae, 127
solid food,
diet, 47
foraging, 47
liquid food, 47
Solenopsis invicta, 47
success rates, 47
Sophora japonica,
China, 602
Cydia trasias, 602
mass trapping, 602
sex pheromone, 602
tortrix, 602
southwestern corn borer,
Diatraea grandiosella, 676
European corn borer, 676
Ostrinia nubilalis, 676
resistance management, 676
simulation modeling, 676
soybean,
alfalfa, 92
Leiobunum, 92
Phalangium opilio, 92
phenology, 92
population, 92
Spalangia cameroni, 411, 425, 505
biological control, 425, 505
Dihrinus himalayanus, 411
dispersal, 425
host age, 505
host size, 505
mark recapture, 425
Musca domestica, 411, 425
Muscidifurax raptor, 411
sex ratio, 505
Spalangia endius, 411
Spalangia gemina, 411
spatial distribution, 425
Spalangia endius,
Dihrinus himalayanus, 411
Musca domestica, 411
Muscidifurax raptor, 411
Spalangia cameroni, 411
Spalangia gemina, 411
Spalangia gemina,
Dihrinus himalayanus, 411
Musca domestica, 411
Muscidifurax raptor, 411
Spalangia cameroni, 411
Spalangia endius, 411
spatial autocorrelation,
geostatistics, 273
moisture, 273
Musca domestica, 273
spatial dynamics, 273
temporal dynamics, 273
spatial distribution, 331, 425
binomial sampling, 331
biological control, 425
Dicyphus hesperus, 331
dispersal, 425
mark recapture, 425
Musca domestica, 425
Spalangia cameroni, 425
spatial dynamics,
geostatistics, 273
moisture, 273
Musca domestica, 273
spatial autocorrelation, 273
temporal dynamics, 273
spatial pattern,
decision rules, 1251
Listronotus oregonensis, 1251
Petroselinum crispum, 1251
probability density functions, 1251
transect pattern, 1251
species richness estimation,
montane forest spiders, 319
sampling intensity, 319
sampling methods, 319
specificity,
allylglucosinolate, 714
Brassicaceae, 714
glucobrassicin, 714
induced defense, 714
spiders,
Carabidae, 785
forest litter arthropods, 785
Oregon, 785
pitfall traps, 785
prescribed fire, 785
Spodoptera exigua, 142, 953, 1197
alfalfa, 953
Bacillus thuringiensis, 1197
beet armyworm, 142
biological control, 1197
biomagnification, 953
Geocoris punctipes, 1197
host plants, 142
Orius tristicolor, 1197
oviposition preference, 142
phytoextraction, 953
phytoremediation, 953
selenium, 953
Zelus renardii, 1197

spread,
Cotesia glomerata, 367
Cotesia rubecula, 367
impact, 367
introduction, 367
Pieris rapae, 367
springtails,
agricultural disturbance, 37
Collembola, 37
cropping systems, 37
soil ecology, 37
stalkborer,
biological control, 432
development rate, 432
Eoreuma loftini, 432
longevity, 432
Lydella jalisco, 432
Staphylindae,
Araneae, 438
biodiversity, 438
Carabidae, 438
pitfall trap size, 438
sampling, 438
Steinernema carpocapsae,
cranberry, 895
Heterorhabditis bacteriophora, 895
Heterorhabditis marelatus, 895
Otiorhynchus sulcatus, 895
strawberry, 895
Steinernema scapterisci,
entomopathogenic nematode, 1226
multiparasitism, 1226
Ormia depleta, 1226
Scapteriscus spp., 1226
tachinid, 1226
Stephanitis pyrioides,
Augmentative release, 1184
Chrysoperla rufilabris, 1184
functional response, 1184
Rhinocapsus vanduzeei, 1184
urban landscape, 1184
Stethorus punctillum,
developmental rate, 177
nonlinear models, 177
predator-prey synchrony, 177
temperature, 177
Tetranychus mcdanieli, 177
storage pest,
Acanthoscelides obtectus, 859
beans, 859
Bruchidae, 859
oviposition behavior, 859
parasitoids, 859
storage system,
biological control, 550
Callosobruchus maculatus, 550
Dinarmus basalis, 550
sulphur-containing compounds, 550
toxicity, 550
stored-grain,
Cryptolestes ferrugineus, 816
mortality, 816
movement, 816
temperature gradient, 816
strain PS149B1,
Bacillus thuringiensis, 208
degradation, 208
model, 208
soil, 208
strawberry,
cranberry, 895
Heterorhabditis bacteriophora, 895
Heterorhabditis marelatus, 895
Otiorhynchus sulcatus, 895
Steinernema carpocapsae, 895
success rates,
diet, 47
foraging, 47
liquid food, 47
Solenopsis invicta, 47
solid food, 47
sugar sprays,
Bacillus thuringiensis, 1231
biological control, 1231
cabbage, 1231
Pieris rapae, 1231
sugarcane,
biological control, 1219
Diatraea saccharalis, 1219
GNA, 1219
nontarget effects, 1219
snowdrop lectin, 1219
sulphur-containing compounds,
biological control, 550
Callosobruchus maculatus, 550
Dinarmus basalis, 550
storage system, 550
toxicity, 550
sunflowers,
crop domestication, 626
Homoeosoma electellum, 626
native agricultural habitats, 626
parasitism, 626
tritrophic interactions, 626
supercooling point,
Anthonomus grandis, 754
boll weevil, 754
surface activity,
behavior, 476
damage, 476
mole crickets, 476
Scapteriscus, 476
soil moisture, 476
survival, 221, 941
development time, 221
energetic reserves, 941
kairomones, 941
nutrition, 941
pharmacophagy, 941
pheromones, 941
Plutella xylostella, 221
rate function, 221
temperature, 221
survivorship, 299, 686
Carposina sasakii, 686
climatic factors, 299
Deois flavopicta, 299
population dynamics, 299, 686
seasonality, 299
sustainable agriculture,
biological control, 84
conservation tillage, 84
entomopathogens, 84
Syrphidae,
biological control, 339
Dasineura oxycoccana, 339
insecticides, 339
parasitoids, 339
Vaccinium, 339

T

Tachinaephagus zealandicus, 375, 732
development rate, 375
feeding treatment, 732
host density, 732
longevity, 375
muscoid flies, 375, 732
parasitoid female age, 732
temperature, 375, 732
tachinid,
entomopathogenic nematode, 1226
multiparasitism, 1226
Ormia depleta, 1226
Scapteriscus spp., 1226
Steinernema scapterisci, 1226
temperature, 177, 221, 354, 375, 509, 608, 732, 1037
biology, 354
Camponotus vicinus, 1037
Coptotermes formosanus, 509
demography, 354
development, 608
development rate, 375
development time, 221
developmental rate, 177
feeding treatment, 732
Glyptotendipes paripes, 608
habitat, 1037
host, 354
host density, 732
life stages, 608
longevity, 375
moisture, 1037
muscoid flies, 375, 732
nest aggregation, 1037
nesting site, 1037
nonlinear models, 177
nuisance midge, 608
parasitoid, 354

parasitoid female age, 732
Plutella xylostella, 221
predator-prey synchrony, 177
rate function, 221
soldier proportion, 509
Stethorus punctillum, 177
survival, 221
Tachinaephagus zealandicus, 375, 732
termite survival, 509
Tetranychus mcdanieli, 177
wood consumption, 509
temperature dependence,
development, 797
leafminers, 797
Liriomyza huidobrensis, 797
Liriomyza trifolii, 797
mortality, 797
temperature effects,
Argentina, 403
biological control, 403
developmental times, 403
fire ant, 403
parasitoid, 403
temperature gradient,
Cryptolestes ferrugineus, 816
mortality, 816
movement, 816
stored-grain, 816
temperature-dependent development,
biological control, 418
brown citrus aphid, 418
Lysiphlebia mirzai, 418
temporal density-dependence,
Apanteles, 988
Goniozus, 988
mode of parasitism, 988
tortricid leafrollers, 988
temporal dynamics,
geostatistics, 273
moisture, 273
Musca domestica, 273
spatial autocorrelation, 273
spatial dynamics, 273
terminal mortality,
Aculops lobuliferus, 1000
defoliation, 1000
intensive forestry, 1000
Populus deltoides, 1000
termite survival,
Coptotermes formosanus, 509
soldier proportion, 509
temperature, 509
wood consumption, 509
termites,
fecundity, 313
nitrogen limitation, 313
ovaries, 313
reproduction, 313
Zootermopsis angusticollis, 313
territoriality,
foraging, 293
inbreeding, 293
social organization, 293
tethered-flight,
age, 1
flight potential, 1
migration, 1
Pseudaletia unipuncta, 1
sex, 1
(Z)-7-tetradecen-2-one,
Exomala orientalis, 1268
highbush blueberries, 1268
mating disruption, 1268
ornamental nurseries, 1268
sex pheromone, 1268
Tetranychidae,
chlorophyll, 215
fluorescence, 215
raspberry, 215
Tetranychus urticae, 215
Tetranychus mcdanieli,
developmental rate, 177
nonlinear models, 177
predator-prey synchrony, 177
Stethorus punctillum, 177
temperature, 177
Tetranychus urticae, 215, 594
carbon dioxide enrichment, 594
chlorophyll, 215
fluorescence, 215
raspberry, 215
Tetranychidae, 215
Trifolium repens, 594
twospotted spider mite, 594
white clover, 594
Thanasimus,
conifer, 1043
Ips, 1043
Platysoma, 1043
population dynamics, 1043
predators, 1043
Thanasimus dubius,
exotic pest, 261
Ips pini, 261
nonindigenous insect, 261
pheromone trapping, 261
predators, 261
Thanasimus undatulus,
Cleridae, 766
Enoclerus sphegeus, 766
Kairomone, 766
Scolytidae, 766
Trypodendron lineatum, 766
Thelohania solenopsae, 127, 887, 1091
biological control, 887, 1091
black imported fire ant, 127
colony size, 1091
dual infections, 887
fire ants, 1091
hybrid fire ant, 127
intracolonial prevalence, 887
microsporidia, 1091
red imported fire ant, 127
Solenopsis invicta, 127, 887
Solenopsis richteri, 127
Vairimorpha invictae, 887
thermal constant,
developmental temperature threshold, 661
egg development, 661
Macrolophus pygmaeus, 661
Miridae, 661
nymphal development, 661
thinning,
Carabidae, 242
forest health, 242
forest management, 242
Pinus ponderosa, 242
prescribed fire, 242
Thripidae,
Allantonematidae, 558
biological control, 558
density dependence, 558
Frankliniella fusca, 558
peanut, 558
thrips,
dendrobium orchids, 523
Frankliniella occidentalis, 523
scouting methods, 523
time constraints,
life history, 702
oviposition, 702
photoperiod, 702
reproductive output, 702
season, 702
time-dependent transition probabilities,
Aedes aegypti, 54
development, 54
food availability, 54
mortality, 54
Plagiorchis elegans, 54
Tiphia,
biological control, 573
gregarinidae, 573
Istocheta, 573
Ovavesicula, 573
Popillia japonica, 573
tomato,
biological control, 397
Colorado potato beetle, 1110
Eretmocerus, 397
life table, 1110
parasitoids, 397
population dynamics, 1110
potatoes, 1110
tritrophic, 397
whitefly, 397
tortricid leafrollers,
Apanteles, 988
Goniozus, 988

mode of parasitism, 988
temporal density-dependence, 988
tortricids,
codling moth, 960
obliquebanded leafroller, 960
oriental fruit moth, 960
pheromone, 960
piezoelectric sprayer, 960
tortrix,
China, 602
Cydia trasias, 602
mass trapping, 602
sex pheromone, 602
Sophora japonica, 602
tospovirus,
brachypterous, 1282
disease management, 1282
Frankliniella fusca, 1282
population dynamics, 1282
TSWV, 1282
toxicity,
biological control, 550
Callosobruchus maculatus, 550
Dinarmus basalis, 550
storage system, 550
sulphur-containing compounds, 550
Toxoptera citricida,
azadirachtin, 172
brown citrus aphid, 172
Lysiphlebus testaceipes, 172
neem extract, 172
Neemix, 172
trait variability,
biomonitoring, 965
developmental instability, 965
environmental stress, 965
Helicoverpa, 965
transect pattern,
decision rules, 1251
Listronotus oregonensis, 1251
Petroselinum crispum, 1251
probability density functions, 1251
spatial pattern, 1251
transgenic,
Bacillus thuringiensis, 1213
beneficial insects, 1213
corn rootworm, 1213
Cry3Bb, 1213
transgenic cotton,
Bacillus thuringiensis, 30
Cry1Ac protein, 30
environmental risk, 30
enzyme-linked immunosorbent assays, 30
insect bioassay, 30
transgenic soybeans,
Collembola, 449
herbicide, 449
microarthropods, 449
population ecology, 449
soil mesofauna, 449
trap,
apple, 1259
Conotrachelus, 1259
peach, 1259
plum curculio, 1259
sampling, 1259
trap catch,
Bark beetles, 982
biological control, 982
vertical distribution, 982
trapping,
coffee pest, 516
Idiarthron subquadratum, 516
mark-recapture, 516
population estimation, 516
sampling, 516
Triaspis aequoris,
biological control, 844
Braconidae, 844
Curculionidae, 844
red sunflower seed weevil, 844
Smicronyx fulvus, 844
Trichogramma,
biological control, 482
life tables, 482
Ostrinia nubilalis, 482
Trichogramma augmentation,
augmentative biological control, 361
cotton, 361
microclimate, 361
microhabitat, 361
Trichogramma evanescens,
connectivity, 1153
Plant structure, 1153
size, 1153
trophic interactions, 1153
Trifolium repens,
carbon dioxide enrichment, 594
Tetranychus urticae, 594
twospotted spider mite, 594
white clover, 594
Trigonospila brevifacies,
biocontrol, 388
foodweb, 388
host competition, 388
leafrollers, 388
shared parasitism, 388
tritrophic,
biological control, 397
Eretmocerus, 397
parasitoids, 397
tomato, 397
whitefly, 397
tritrophic interactions, 626, 874
body size, 874
Chrysocharis flacilla, 874
crop domestication, 626
Halticoptera helioponi, 874
Homoeosoma electellum, 626
Liriomyza huidobrensis, 874
native agricultural habitats, 626
parasitism, 626
Phaedrotoma scabriventris, 874
sunflowers, 626
trophic interactions,
connectivity, 1153
Plant structure, 1153
size, 1153
Trichogramma evanescens, 1153
Trypodendron lineatum,
Cleridae, 766
Enoclerus sphegeus, 766
Kairomone, 766
Scolytidae, 766
Thanasimus undatulus, 766
TSWV,
brachypterous, 1282
disease management, 1282
Frankliniella fusca, 1282
population dynamics, 1282
tospovirus, 1282
turfgrass,
conservation biological control, 564
ornamentals, 564
parasitoids, 564
predators, 564
wildflowers, 564
twospotted spider mite,
carbon dioxide enrichment, 594
Tetranychus urticae, 594
Trifolium repens, 594
white clover, 594

U

urban landscape,
Augmentative release, 1184
Chrysoperla rufilabris, 1184
functional response, 1184
Rhinocapsus vanduzeei, 1184
Stephanitis pyrioides, 1184

V

Vaccinium, 339, 852
Acrobasis, 852
biological control, 339
blueberry, 852
cherry fruitworm, 852
cranberry fruitworm, 852
Dasineura oxycoccana, 339
Grapholita, 852
insecticides, 339
parasitoids, 339
Syrphidae, 339
Vairimorpha invictae,
biological control, 887
dual infections, 887

intracolonial prevalence, 887
Solenopsis invicta, 887
Thelohania solenopsae, 887
vegetable,
Bemisia tabaci, 1030
biocontrol, 1030
Encarsia, 1030
Eretmocerus, 1030
whitefly, 1030
vegetation,
Anopheles, 134
breeding, 134
mosquito larvae, 134
water velocity, 134
vertical distribution,
Bark beetles, 982
biological control, 982
trap catch, 982
vineyards,
domatia, 648
Kampimodromus aberrans, 648
leaf trichomes, 648
pollen, 648
voltinism,
bruchid, 469
Gleditsia japonica, 469
herbivorous insect, 469
overwintering stage, 469
seasonal adaptation, 469

W

walnuts,
California, 723
Cydia pomonella, 723
nontarget impacts, 723
parasitoid, 723
water velocity,
Anopheles, 134
breeding, 134
mosquito larvae, 134
vegetation, 134
Weibull function,
adult emergence, 864
biofix, 864
corn rootworm, 864
Diabrotica, 864
western spruce budworm,
Bacillus thuringiensis subsp. *kurstaki*, 903
Choristoneura occidentalis, 903
Douglas-fir, 903
nontarget Lepidoptera, 903
white clover,
carbon dioxide enrichment, 594
Tetranychus urticae, 594
Trifolium repens, 594
twospotted spider mite, 594
white pine weevil,
interior spruce, 198
Mexican spruce, 198
Pissodes strobi, 198
Sitka spruce, 198
whitefly, 107, 397, 1030
Abutilon theophrasti, 107
Acalypha ostryaefolia, 107
Bemisia tabaci, 1030
biocontrol, 1030
biological control, 397
Coleomegilla maculata, 107
Encarsia, 1030
Eretmocerus, 397, 1030
oviposition, 107
parasitoids, 397
tomato, 397
tritrophic, 397
vegetable, 1030
Zea mays, 107
whitemarked tussock moth,
Betula papyrifera, 10
condensed tannin, 10
Orgyia leucostigma, 10
paper birch, 10
wildflowers,
conservation biological control, 564
ornamentals, 564
parasitoids, 564
predators, 564
turfgrass, 564
wind,
Anthonomus grandis, 1128
boll weevil, 1128
flight behavior, 1128
pheromone traps, 1128
sampling, 1128
wood consumption,
Coptotermes formosanus, 509
soldier proportion, 509
temperature, 509
termite survival, 509
woody plants,
fertilization, 923
growth/differentiation balance, 923
host plant resistance, 923
secondary metabolism, 923

Z

Zea mays, 107, 112
Abutilon theophrasti, 107, 112
Acalypha ostryaefolia, 107, 112
Coleomegilla maculata, 107, 112
oviposition, 107, 112
predation, 112
whitefly, 107
Zelus renardii,
Bacillus thuringiensis, 1197
biological control, 1197
Geocoris punctipes, 1197
Orius tristicolor, 1197
Spodoptera exigua, 1197
Zootermopsis angusticollis,
fecundity, 313
nitrogen limitation, 313
ovaries, 313
reproduction, 313
termites, 313